

WO02094787

Publication Title:

2-OXO-PIPERIDINYL- AND 2-OXO-AZEPANYL ALKANOIC ACID DERIVATIVES FOR THE TREATMENT OF EPILEPSY AND OTHER NEUROLOGICAL DISORDERS

Abstract:

Abstract of WO02094787

This invention relates to new 2-oxo-piperidinyl- and 2-oxo-azepanyl alkanolic acid derivatives of formula (1) having anticonvulsant activity which are useful as therapeutic agents for the treatment or prevention of epilepsy and other neurological disorders. The invention also concerns processes for preparing these derivatives and novel intermediates used in the preparation of these derivatives. Data supplied from the esp@cenet database - Worldwide

Courtesy of <http://v3.espacenet.com>

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 November 2002 (28.11.2002)

PCT

(10) International Publication Number
WO 02/094787 A1

(51) International Patent Classification⁷: **C07D 211/76**,
211/78, 409/04, 223/10, A61K 31/45, 31/55, A61P 25/08

(21) International Application Number: PCT/EP02/05503

(22) International Filing Date: 17 May 2002 (17.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01112541.6 23 May 2001 (23.05.2001) EP

(71) Applicant (for all designated States except US): **UCB**,
S.A. [BE/BE]; Allée de la Recherche, 60, B-1070 Bruxelles (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MICHEL, Philippe**
[BE/BE]; Stoofstraat, 15, B-1650 BEERSEL (BE).
KENDA, Benoît [BE/BE]; Trieux des frênes 5, B-5080
EMINES (BE).

(74) Agent: **LECHIEN, Monique**; UCB, S.A., Intellectual
Property Department, Allée de la Recherche, 60, B-1070
Bruxelles (BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 2-OXO-PIPERIDINYL- AND 2-OXO-AZEPANYL ALKANOIC ACID DERIVATIVES FOR THE TREATMENT OF EPILEPSY AND OTHER NEUROLOGICAL DISORDERS



(57) Abstract: This invention relates to new 2-oxo-piperidinyl- and 2-oxo-azepanyl alkanolic acid derivatives of formula (1) having anticonvulsant activity which are useful as therapeutic agents for the treatment or prevention of epilepsy and other neurological disorders. The invention also concerns processes for preparing these derivatives and novel intermediates used in the preparation of these derivatives.

WO 02/094787 A1

2-OXO-PIPERIDINYL- AND 2-OXO-AZEPANYL ALKANOIC ACID DERIVATIVES
FOR THE TREATMENT OF EPILEPSY AND OTHER NEUROLOGICAL DISORDERS

5 FIELD OF THE INVENTION

This invention relates to new 2-oxo-piperidinyl- and 2-oxo-azepanyl alkanolic acid derivatives having anticonvulsant activity which are useful as therapeutic agents for the treatment and/or prevention of epilepsy and other neurological disorders. The invention also
10 concerns processes for preparing these derivatives and novel intermediates used in the preparation of these derivatives.

BACKGROUND OF THE INVENTION

15 Epilepsy is a relatively common neurological condition affecting 0.4-1 % of the world's population (45-100 million people). For the general population there are approximately 20-70 new cases per 100,000 diagnosed each year with a 3-5 % lifetime probability of developing the disease. The older established antiepileptic drugs (AEDs) phenytoin, carbamazepine, clonazepam, ethosuximide, valproic acid and barbiturates are widely prescribed but suffer from
20 a range of side effects. Furthermore, there is a significant group of patients (20-30 %) that are resistant to the currently available therapeutic agents. Since 1989 several new drugs have been launched, including felbamate, gabapentin, lamotrigine, oxcarbazepine, tiagabine, topiramate, vigabatrin, zonisamide and levetiracetam. While many of the new AEDs show improved efficacies and side-effect profiles, patients with intractable epilepsy remain untreated. There is clearly a
25 need for improved medications (Cosford N. D. P. et al., Annual Reports in Medicinal Chemistry (1998), 33, 61-70).

US Patent No. 6,066,666 reports lactam and thiolactam derivatives having useful anticonvulsant and anxiolytic activity, and thus being effective in the treatment of epilepsy.

US Patent No. 5,334,720 relates to diphenyl-1-(aminoalkyl)-2-piperidone and -2-
30 pyrrolidinone derivatives, which possess anticonvulsant properties and are useful anti-epileptic agents.

European Patent No. 0 162 036 B1 discloses levetiracetam or (S)-(-)- α -ethyl-2-oxo-1-pyrrolidineacetamide as well as its use as a protective agent for the treatment and the

prevention of hypoxic and ischemic type aggressions of the central nervous system. This compound is also effective in the treatment of epilepsy, a therapeutic indication for which it has been demonstrated that its dextrorotatory enantiomer (R)-(+)- α -ethyl-2-oxo-1-pyrrolidineacetamide completely lacks activity (Gower A.J. et al., Eur. J. Pharmacol. (1992), 222, 193-203). In the European Patent No. 0 645 139 B1, levetiracetam has been disclosed for its anxiolytic activity. International Patent Application No. PCT/EP00/11808 discloses the use of levetiracetam for the curative and/or prophylactic treatment of bipolar disorders, migraine, chronic or neuropathic pain as well as combinations of levetiracetam with at least one compound inducing neural inhibition mediated by GABA_A receptors.


Continuing its research work in the field of the treatment of epilepsy, the applicant discloses now new 2-oxo-piperidiny- and 2-oxo-azepanyl alkanoic acid derivatives. These new 2-oxo-piperidiny- and 2-oxo-azepanyl alkanoic acid derivatives are useful for the prevention or the treatment of epilepsy, epileptogenesis, seizure disorders and convulsions.

These compounds may also be used for the treatment of other neurological disorders including bipolar disorders, mania, depression, anxiety, migraine, trigeminal and other neuralgia, chronic pain, neuropathic pain, cerebral ischemia, cardiac arrhythmia, myotonia, cocaine abuse, stroke, myoclonus, essential tremor and other movement disorders, neonatal cerebral haemorrhage, amyotrophic lateral sclerosis, spasticity, Parkinson's disease and other degenerative diseases.

In addition the compounds according to the invention may be used in the treatment of bronchial asthma, asthmatic status and allergic bronchitis, asthmatic syndrome, bronchial hyperreactivity and bronchospastic syndromes as well as allergic and vasomotor rhinitis and rhinoconjunctivitis.

DESCRIPTION OF THE INVENTION

This invention provides novel compounds of the formula I



wherein

n represents 0 or 1 whereby R^1 is not existent when $n=0$ and R^1 is existent when $n=1$;

A^1 represents an oxygen or a sulfur atom;

5 X is $-\text{CONR}^7\text{R}^8$, $-\text{COOR}^9$, $-\text{CO-R}^{10}$ or CN;

R^1 when existent, R^2 , R^3 , R^4 and R^5 are the same or different and each is independently hydrogen, halogen, hydroxy, thiol, amino, nitro, nitrooxy, cyano, azido, carboxy, amido, sulfonic acid, sulfonamide, alkyl, alkenyl, alkynyl, ester, ether, aryl, heterocycle, or an oxy derivative, thio derivative, amino derivative, acyl derivative, sulfonyl derivative or sulfinyl derivative,

10 provided that at least one of the substituents R chosen from R^1 when existent, R^2 , R^3 , R^4 or R^5 is not hydrogen;

R^6 is hydrogen, alkyl, aryl or $-\text{CH}_2\text{-R}^{6a}$ wherein R^{6a} is aryl, heterocycle, halogen, hydroxy, amino, nitro or cyano;

15 R^7 , R^8 and R^9 are the same or different and each is independently hydrogen, hydroxy, alkyl, aryl, heterocycle or an oxy derivative; and

R^{10} is hydrogen, hydroxy, thiol, halogen, alkyl, aryl, heterocycle or a thio derivative;

their pharmaceutically acceptable salts, geometrical isomers (including cis and trans, Z and E isomers), enantiomers, diastereoisomers and mixtures thereof (including all possible mixtures of stereoisomers).

20 In the above formula, at least one substituent R^1 to R^5 is different from hydrogen. Some non-substituted compounds are disclosed in US Patent No. 5,468,733 and 5,516,759. US Patent No. 5,468,733 discloses non-ring substituted 2-oxo-1-pyrrolidiny and 2-oxo-1-piperidiny derivatives as inhibitors of the oncogene Ras protein. In particular, these compounds block the ability of Ras to transform normal cells to cancer cells, and therefore can be included in several
25 chemotherapeutic compositions for treating cancer.

US Patent No. 5,516,759 discloses non-ring substituted 2-oxo-1-pyrrolidiny, 2-oxo-1-piperidiny and azepany derivatives present at the N-terminus of dodecapeptides possessing LHRH (luteinizing hormone-releasing hormone) antagonistic activity. Such LHRH antagonists are useful in the treatment of a variety of conditions in which suppression of sex steroids plays a key
30 role including contraception, delay of puberty, treatment of benign prostatic hyperplasia a.o.

In the definitions set forth below, unless otherwise stated, R^{11} and R^{12} are the same or different and each is independently amido, alkyl, alkenyl, alkynyl, acyl, ester, ether, aryl, aralkyl, heterocycle or an oxy derivative, thio derivative, acyl derivative, amino derivative, sulfonyl derivative, or sulfinyl derivative, each optionally substituted with any suitable group, including,
35 but not limited to, one or more moieties selected from lower alkyl or other groups as described below as substituents for alkyl.

The term "oxy derivative", as used herein, is defined as including -O-R¹¹ groups wherein R¹¹ is as defined above except for "oxy derivative". Non-limiting examples are alkoxy, alkenyloxy, alkynyloxy, acyloxy, oxyester, oxyamido, alkylsulfonyloxy, alkylsulfinyloxy, arylsulfonyloxy, arylsulfinyloxy, aryloxy, aralkoxy or heterocycloxy such as pentyloxy, allyloxy, methoxy, ethoxy, phenoxy, benzyloxy, 2-naphthyloxy, 2-pyridyloxy, methylenedioxy, carbonate.

The term "thio derivative", as used herein, is defined as including -S-R¹¹ groups wherein R¹¹ is as defined above except for "thio derivative". Non-limiting examples are alkylthio, alkenylthio, alkynylthio and arylthio.

The term "amino derivative", as used herein, is defined as including -NHR¹¹ or -NR¹¹R¹² groups wherein R¹¹ and R¹² are as defined above. Non-limiting examples are mono- or di-alkyl-, alkenyl-, alkynyl- and arylamino or mixed amino.

The term "acyl derivative", as used herein, represents a radical derived from carboxylic acid and thus is defined as including groups of the formula R¹¹-CO-, wherein R¹¹ is as defined above and may also be hydrogen. Preferred are acyl derivatives of formula -COR¹¹ wherein R¹¹ is selected from hydrogen, C1-12 alkyl, C2-12 alkenyl, C2-12 alkynyl, heterocycle and aryl. Non-limiting examples are formyl, acetyl, propionyl, isobutyryl, valeryl, lauroyl, heptanedioyl, cyclohexanecarbonyl, crotonoyl, fumaroyl, acryloyl, benzoyl, naphthoyl, furoyl, nicotinoyl, 4-carboxybutanoyl, oxalyl, ethoxalyl, cysteinyl, oxamoyl.

The term "sulfonyl derivative", as used herein, is defined as including a group of the formula -SO₂-R¹¹, wherein R¹¹ is as defined above except for "sulfonyl derivative". Non-limiting examples are alkylsulfonyl, alkenylsulfonyl, alkynylsulfonyl and arylsulfonyl.

The term "sulfinyl derivative", as used herein, is defined as including a group of the formula -SO-R¹¹, wherein R¹¹ is as defined above except for "sulfinyl derivative". Non-limiting examples are alkylsulfinyl, alkenylsulfinyl, alkynylsulfinyl and arylsulfinyl.

The term "alkyl", as used herein, is defined as including saturated, monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof and generally containing 1-20 carbon atoms, most often 1 to 12 carbon atoms, preferably 1-7 carbon atoms for non-cyclic alkyl and 3-7 carbon atoms for cycloalkyl (in these two preferred cases, unless otherwise specified, "lower alkyl"), each optionally substituted by, preferably 1 to 5, substituents independently selected from the group consisting of halogen, hydroxy, thiol, amino, nitro, cyano, thiocyanato, acyl, acyloxy, sulfonyl derivative, sulfinyl derivative, alkylamino, carboxy, ester, ether, amido, azido, cycloalkyl, sulfonic acid, sulfonamide, thio derivative, alkylthio, oxyester, oxyamido, heterocycle, vinyl, alkoxy (preferably C1-5), aryloxy (preferably C6-10) and aryl (preferably C6-10).

Preferred are alkyl groups containing 1 to 7 carbon atoms, each optionally substituted by one or more substituents selected from hydroxy, halogen, cyano, thiocyanato, alkoxy, azido,

alkylthio, cyclopropyl, acyl and phenyl. Most preferred are C1-4 alkyl and C3-7 cycloalkyl, each optionally substituted by one or more hydroxy, halogen, lower alkyl or/and azido.

Most preferred alkyl groups are hydroxymethyl, propyl, butyl, 2,2,2-trifluoroethyl, 2-bromo-2,2-difluoroethyl, 2-chloro-2,2-difluoroethyl, 3,3,3-trifluoropropyl, cyclopropylmethyl, iodomethyl, azidomethyl, 2,2-difluoropropyl, 2-iodo-2,2-difluoroethyl.

The term "lower alkyl", as used herein, and unless otherwise specified, refers to C₁ to C₇ saturated straight, branched or cyclic hydrocarbon. Non limiting examples are methyl, ethyl, propyl, isopropyl, butyl, tertibutyl, pentyl, cyclopropyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, 3-methylpentyl, 2,2-dimethylbutyl, optionally substituted with any suitable group, including but not limited to one or more moieties selected from groups as described above for the alkyl groups. Preferably, lower alkyl is methyl.

The term "alkenyl", as used herein, is defined as including both branched and unbranched, unsaturated hydrocarbon radicals having at least one double bond, and being optionally substituted by at least one substituent selected from the group consisting of halogen, hydroxy, thiol, amino, thiocyanato, azido, alkylthio, cycloalkyl, acyl, nitro, cyano, aryl and heterocycle.

Preferred alkenyl groups are C2-C12 alkenyls, especially C2-6 alkenyls, such as ethenyl (= vinyl), 1-methyl-1-ethenyl, 2,2-dimethyl-1-ethenyl, 1-propenyl, 2-propenyl (= allyl), 1-butenyl, 2-butenyl, 3-butenyl, 4-pentenyl, 1-methyl-4-pentenyl, 3-methyl-1-pentenyl, 1-hexenyl, 2-hexenyl and the like, optionally being substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, cycloalkyl, phenyl and acyl. Most preferred is vinyl, optionally substituted by one or more halogen or/and lower alkyl, and especially 2,2-difluorovinyl, 2,2-dibromovinyl and 2,2-dichlorovinyl.

The term "alkynyl" as used herein, is defined as including a monovalent branched or unbranched hydrocarbon radical containing at least one carbon-carbon triple bond, for example ethynyl, 2-propynyl (= propargyl), and the like, and being optionally substituted by at least one substituent selected from the group consisting of halogen, hydroxy, thiol, amino, nitro, cyano, aryl, heterocycle, thiocyanato, azido, alkylthio, alkyl and acyl.

Preferred alkynyl groups are C2-12 alkynyl, especially C2-6 alkynyl, optionally being substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, acyl, aryl such as phenyl and alkyl, preferably cycloalkyl.

Most preferred are ethynyl, propynyl and butynyl, optionally substituted by lower alkyl or/and halogen, and especially 1-propynyl, cyclopropylethynyl, 3-methyl-1-butynyl and 3,3,3-trifluoro-1-propynyl.

When present as bridging groups, alkyl, alkenyl and alkynyl represent straight- or branched chains, C1-12, preferably C1-4-alkylene or C2-12-, preferably C2-4-alkenylene or -alkynylene moieties respectively.

Groups where branched derivatives are conventionally qualified by prefixes such as "n", "sec", "iso" and the like (e.g. "n-propyl", "sec-butyl") are in the n-form unless otherwise stated.

The term "aryl", as used herein, is defined as including an organic radical derived from an aromatic hydrocarbon consisting of at least one ring, most often 1 to 3 rings and generally containing 6-30 carbon atoms by removal of one hydrogen, such as phenyl and naphthyl, each optionally substituted by one or more substituents independently selected from halogen, hydroxy, thiol, amino, nitro, cyano, acyl, acyloxy, sulfonyl, sulfinyl, alkylamino, carboxy, ester, ether, amido, azido, sulfonic acid, sulfonamide, alkylsulfonyl, alkylsulfinyl, C1-6-alkylthio, oxyester, oxyamido, aryl, C1-6-alkoxy, C6-10-aryloxy, C1-6-alkyl, C1-6-haloalkyl. Aryl radicals are preferably monocyclic or bicyclic containing 6-10 carbon atoms. Preferred aryl groups are phenyl and naphthyl each optionally substituted by one or more substituents independently selected from halogen, nitro, amino, azido, C1-6-alkoxy, C1-6-alkyl, C1-6-haloalkyl, sulfonyl and phenyl.

Preferred aryl is phenyl, optionally substituted by one or more halogen, lower alkyl, azido or nitro, such as 3-chlorophenyl and 3-azidophenyl.

The term "halogen", as used herein, includes an atom of Cl, Br, F, I.

The term "hydroxy", as used herein, represents a group of the formula -OH.

The term "thiol", as used herein, represents a group of the formula -SH.

The term "cyano", as used herein, represents a group of the formula -CN.

The term "nitro", as used herein, represents a group of the formula -NO₂.

The term "nitrooxy", as used herein, represents a group of the formula -ONO₂.

The term "amino", as used herein, represents a group of the formula -NH₂.

The term "azido", as used herein, represents a group of the formula -N₃.

The term "carboxy", as used herein, represents a group of the formula -COOH.

The term "sulfonic acid", as used herein, represents a group of the formula -SO₃H.

The term "sulfonamide", as used herein, represents a group of the formula -SO₂NH₂.

The term "ester", as used herein, is defined as including a group of formula -COO-R¹¹ wherein R¹¹ is as defined above except oxy derivative, thio derivative or amino derivative. Preferred are esters of formula -COOR¹¹ wherein R¹¹ is selected from C1-12 alkyl, C2-12 alkenyl, C2-12 alkynyl and aryl. Most preferred are esters where R¹¹ is a lower alkyl, especially methyl.

The term "ether" is defined as including a group selected from C1-50- straight or branched alkyl, or C2-50- straight or branched alkenyl or alkynyl groups or a combination of the

same, interrupted by one or more oxygen atoms.

The term "amido" is defined as including a group of formula $-\text{CONH}_2$ or $-\text{CONHR}^{11}$ or $-\text{CONR}^{11}\text{R}^{12}$ wherein R^{11} and R^{12} are as defined above.

The term "heterocycle", as used herein, is defined as including an aromatic or non
5 aromatic cyclic alkyl, alkenyl, or alkynyl moiety as defined above, having at least one O, S
and/or N atom interrupting the carbocyclic ring structure and optionally, one of the carbon of
the carbocyclic ring structure may be replaced by a carbonyl, and optionally being substituted
with any suitable group, including but not limited to one or more moieties selected from lower
alkyl, or other groups as described above for the alkyl groups. Non-limiting examples of
10 heterocycles are pyridyl, furyl, pyrrolyl, thienyl, isothiazolyl, triazolyl, imidazolyl, benzimidazolyl,
tetrazolyl, quinazolinyl, quinoliziny, naphthyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, quinolyl,
isoquinolyl, isobenzofuranyl, benzothienyl, pyrazolyl, indolyl, indoliziny, purinyl, isoindolyl,
carbazolyl, thiazolyl, 1,2,4-thiadiazolyl, thiomorpholinyl, thieno(2,3-b)furanly, furopyranly,
benzofuranly, benzoxepinyl, isooxazolyl, oxazolyl, thianthrenyl, benzothiazolyl, or benzoxazolyl,
15 cinnolinyl, phthalazinyl, quinoxalinyl, 1-oxidopyridyl, phenanthridinyl, acridinyl, perimidinyl,
phenanthrolinyl, phenothiazinyl, furazanyl, benzodioxolyl, isochromanly, indolinyl, xanthenyl,
hypoxanthinyl, pteridinyl, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl,
pyrrolopyrimidinyl, pyrazolopyrimidinyl, tetrahydrofuranly, tetrahydropyranly, piperidinyl,
piperidyl, piperazinyl, imidazolidinyl, morpholino, morpholinyl, 1-oxaspiro(4.5)dec-2-yl,
20 pyrrolidinyl, 2-oxo-pyrrolidinyl, sugar moieties (i.e. glucose, pentose, hexose, ribose, fructose,
which may also be substituted) optionally substituted by alkyl or as described above for the alkyl
groups. The term "heterocycle" also includes bicyclic, tricyclic and tetracyclic, spiro groups in
which any of the above heterocyclic rings is fused to one or two rings independently selected
from an aryl ring, a cyclohexane ring, a cyclohexene ring, a cyclopentane ring, a cyclopentene
25 ring or another monocyclic heterocyclic ring or where a monocyclic heterocyclic group is bridged
by an alkylene group, such as quinuclidinyl, 7-azabicyclo(2.2.1)heptanyl, 7-
oxabicyclo(2.2.1)heptanyl, 8-azabicyclo(3.2.1)octanyl.

The heterocycle is preferably selected from triazolyl, tetrazolyl, pyrrolidinyl, pyridyl, 1-
oxidopyridyl, thiomorpholinyl, benzodioxolyl, furyl, oxazolyl, pyrimidinyl, pyrrolyl, thiadiazolyl,
30 thiazolyl, thienyl and piperazinyl, each optionally substituted by one or more substituents
selected from halogen, alkyl, substituted alkyl, alkoxy, nitro, amino, acyl and phenyl.

More preferably the heterocycle is selected from tetrazolyl, pyrrolidinyl, pyridyl, furyl, pyrrolyl,
thiazolyl and thienyl, each optionally substituted by one or more substituents selected from
halogen, alkyl, halogen substituted alkyl, acyl, alkoxy, nitro, amino and phenyl, and especially
35 from 2-and 3-thienyl, optionally substituted by one or more halogen, acyl such as formyl, cyano
and/or lower alkyl, such as methyl.

In the above definitions it is to be understood that when a substituent such as R¹, R², R³, R⁴, R⁵, R⁷, R⁸, R⁹, R¹⁰ is attached to the rest of the molecule via a heteroatom or a carbonyl, a straight- or branched chain, C1-12-, preferably C1-4-alkylene or C2-12, preferably C2-4-alkenylene or -alkynylene bridge may optionally be interposed between the heteroatom or the carbonyl and the point of attachment to the rest of the molecule.

The "pharmaceutically acceptable salts" according to the invention include therapeutically active, non-toxic base and acid salt forms which the compounds of formula (I) are able to form.

The acid addition salt form of a compound of formula (I) that occurs in its free form as a base can be obtained by treating said free base form with an appropriate acid such as an inorganic acid, for example, a hydrohalic such as hydrochloric or hydrobromic, sulfuric, nitric, phosphoric and the like; or an organic acid, such as, for example, acetic, hydroxyacetic, propanoic, lactic, pyruvic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, p-toluenesulfonic, cyclamic, salicylic, p-aminosalicylic, pamoic and the like.

The compounds of formula (I) containing acidic protons may be converted into their therapeutically active, non-toxic base addition salt form, e.g. metal or amine salts, by treatment with appropriate organic and inorganic bases. Appropriate base salt forms include, for example, ammonium salts, alkali and earth alkaline metal salts, e.g. lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. N-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

Conversely said salt forms can be converted into the free forms by treatment with an appropriate base or acid.

Compounds of the formula I and their salts can be in the form of a solvate, which is included within the scope of the present invention. Such solvates include for example hydrates, alcoholates and the like.

Many of the compounds of formula I and some of their intermediates have at least one stereogenic center in their structure. This stereogenic center may be present in a R or a S configuration, said R and S notation is used in correspondence with the rules described in Pure Appl. Chem. (1976), 45, 11-30.

The invention also relates to all stereoisomeric forms such as enantiomeric and diastereoisomeric forms of the compounds of formula I or mixtures thereof (including all possible mixtures of stereoisomers).

Furthermore, certain compounds of formula I which contain alkenyl groups may exist as Z (zusammen) or E (entgegen) isomers. In each instance, the invention includes both mixture and separate individual isomers.

Multiple substituents on the piperidinyl or the azepanyl ring can also stand in either cis or trans relationship to each other with respect to the plane of the piperidinyl or the azepanyl ring.

Some of the compounds of formula I may also exist in tautomeric forms. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.


With respect to the present invention reference to a compound or compounds is intended to encompass that compound in each of its possible isomeric forms and mixtures thereof unless the particular isomeric form is referred to specifically.

The invention also includes within its scope prodrug forms of the compounds of formula I and its various sub-scopes and sub-groups.


The term "prodrug" as used herein includes compound forms which are rapidly transformed in vivo to the parent compound according to the invention, for example, by hydrolysis in blood. Prodrugs are compounds bearing groups which are modified by biotransformation prior to exhibiting their pharmacological action. Such groups include moieties which are readily oxidised, cyclised or cleaved, which compound after biotransformation remains or becomes pharmacologically active. For example, metabolically cleavable groups form a class of groups well known to practitioners of the art. They include, but are not limited to such groups as alkanoyl (i.e. acetyl, propionyl, butyryl, and the like), unsubstituted and substituted carbocyclic aroyl (such as benzoyl, substituted benzoyl and 1- and 2-naphthoyl), alkoxycarbonyl (such as ethoxycarbonyl), trialkylsilyl (such as trimethyl- and triethylsilyl), monoesters formed with dicarboxylic acids (such as succinyl), phosphate, sulfate, sulfonate, sulfonyl, sulfinyl and the like. The compounds bearing the biotransformable groups have the advantage that they may exhibit improved bioavailability as a result of enhanced solubility and/or rate of absorption conferred upon the parent compound by virtue of the presence of the biotransformable group. T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery System", Vol. 14 of the A.C.S. Symposium Series; "Bioreversible Carriers in Drug Design", ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

The term "R substituent" refers to R^1 , R^2 , R^3 , R^4 or R^5 , independently.

According to a preferred embodiment, the present invention relates to a compound of formula I as defined above wherein n represents 0. The compound is a 6-ring structure (2-thioxo- or 2-oxo-piperidinyl derivative) wherein R^1 is not existent since $n=0$, and is depicted by the formula (I-A).



According to a following embodiment, the present invention relates to a compound of formula I according to the invention as defined above wherein n represents 1. The compound is a 7-ring structure (2-thioxo- or 2-oxo-azepanyl derivative) wherein R¹ is existent since n=1 and depicted by the formula (I-B).



According to a more preferred embodiment, the invention relates to said compound as defined above wherein n=0, R³ and/or R⁴ are different from hydrogen and R² and R⁵ represent hydrogen.

According to another more preferred embodiment, the invention relates to said compound as defined above wherein n=1, R², R³ and/or R⁴ are different from hydrogen and wherein R¹ and R⁵ represent hydrogen.

According to a yet more preferred embodiment, the invention relates to said compound as defined above wherein only one R substituent chosen from R³ or R⁴ when n=0 or from R², R³ or R⁴ when n=1, is different from hydrogen and the remaining R substituent(s) is/are hydrogen. We hereby refer to a mono-substituted 2-thioxo- or 2-oxo-piperidinyll or 2-thioxo- or 2-oxo-azepanyl derivatives.

According to another preferred embodiment, the present invention relates to compounds of formula I according to the invention as defined above wherein A¹ represents an oxygen atom. We hereby refer to 2-oxo-piperidinyll or 2-oxo-azepanyl derivatives.

According to another preferred embodiment, the present invention relates to compounds of formula I according to the invention as defined above wherein X is CONR⁷R⁸, especially

CONH₂. We hereby refer to amido derivatives of 2-oxo(or thioxo)-piperidinyl or 2-oxo(or thioxo)-azepanyl.

According to another preferred embodiment, the present invention relates to compounds of formula I according to the invention as defined above wherein R⁶ represents hydrogen, C1-4 alkyl, or a CH₂-R^{6a} group wherein R^{6a} represents a heterocycle. Most preferably R⁶ is a C1-4 alkyl, especially ethyl. When R⁶ is ethyl we refer to 2-(2-oxo(or thioxo)-1-piperidinyl)butanamide or 2-(2-oxo(or thioxo)-1-azepanyl)butanamide derivatives.

According to another preferred embodiment, the present invention relates to compounds of formula I according to the invention as defined above wherein the carbon atom to which R⁶ is attached is of the S configuration. In case where R⁶ is ethyl, A is oxygen and X is CON R⁷ R⁸, we refer then to (2S)-2-(2-oxo-1-piperidinyl)butanamide or (2S)-2-(2-oxo-1-azepanyl)butanamide derivatives.

According to a preferred embodiment, the present invention relates to a compound as defined above wherein R² when n=1, R³ and R⁴ are the same or different and each is independently hydrogen, halogen, nitro, nitrooxy, cyano, carboxy, amido, sulfonic acid, sulfonamide, alkyl, alkenyl, alkynyl, ester, ether, aryl, heterocycle, acyl derivative, sulfonyl derivative or sulfinyl derivative;

R¹ when existent, R² when n=0 and R⁵ are hydrogen;

R⁶ is hydrogen, alkyl, aryl or -CH₂-R^{6a} wherein R^{6a} is aryl, heterocycle, halogen,

hydroxy, amino, nitro or cyano;

provided that, when R⁶ is hydrogen, X is -CONR⁷R⁸ and that the compound is neither methyl (2R)-2-[(6R)-6-methyl-2-oxoazepanyl]-3-phenylpropanoate nor methyl (2S)-2-[(4R)-4-methyl-2-oxoazepanyl]-3-phenylpropanoate.

According to this preferred embodiment, the compound is generally such that when R⁶ is benzyl, X is -COOCH₃ and n=1, R² is different from methyl when R³ and R⁴ are both hydrogen and R⁴ is different from methyl when R² and R³ are both hydrogen.

According to another preferred embodiment, the present invention relates to a compound as defined above wherein R² when n=1, R³ and R⁴ are the same or different and each is independently hydrogen; cyano; carboxy; amido;

C1-12 alkyl, each optionally substituted by one or more substituents selected from hydroxy, halogen, cyano, thiocyanato, alkoxy, azido, alkylthio, cycloalkyl, acyl, aryl and heterocycle;

C2-12 alkenyl, each optionally substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, alkyl, aryl and acyl;

C2-12 alkynyl, each optionally substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, alkyl, aryl and acyl;

acyl derivative of formula $-\text{CO}-\text{R}^{11}$, wherein R^{11} is selected from C1-12 alkyl, C2-12 alkenyl, C2-12 alkynyl, heterocycle and aryl;

ester of formula $-\text{CO}-\text{O}-\text{R}^{11}$ wherein R^{11} is selected from C1-12 alkyl, C2-12 alkenyl, C2-12 alkynyl and aryl;

- 5 heterocycle selected from triazolyl, tetrazolyl, pyrrolidinyl, pyridyl, 1-oxidopyridyl, thiomorpholinyl, benzodioxolyl, furyl, oxazolyl, pyrimidinyl, pyrrolyl, thiadiazolyl, thiazolyl, thienyl and piperazinyl, each optionally substituted by one or more substituents selected from halogen, alkyl, substituted alkyl, alkoxy, nitro, amino, acyl and phenyl; aryl, each optionally substituted by one or more substituents selected from C1-6 alkyl, C1-6
10 haloalkyl, C1-6 alkoxy, C1-6 alkylthio, amino, azido, sulfonyl, aryl and nitro.

According to another preferred embodiment, the present invention relates to a compound as defined above, wherein R^2 when $n=1$, R^3 and R^4 are the same or different and each is independently hydrogen;

- C1-7 alkyl, each optionally substituted by one or more substituents selected from hydroxy,
15 halogen, cyano, thiocyanato, alkoxy, azido, alkylthio, cyclopropyl, acyl and phenyl;
C2-6 alkenyl, each optionally substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, cycloalkyl, phenyl and acyl;
C2-6 alkynyl, each optionally substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, cycloalkyl, phenyl and acyl;
20 heterocycle selected from tetrazolyl, pyrrolidinyl, pyridyl, furyl, pyrrolyl, thiazolyl and thienyl, each optionally substituted by one or more substituents selected from halogen, alkyl, halogen substituted alkyl, acyl, alkoxy, nitro, amino and phenyl;
phenyl, each optionally substituted by one or more substituents selected from C1-6 alkyl, halogen substituted alkyl, halogen, alkoxy, amino, azido, sulfonyl, phenyl and nitro.

- 25 According to another preferred embodiment, the present invention relates to a compound as defined above wherein at least one of the R substituents chosen from the group R^2 , R^3 and R^4 when $n=1$ or from the group R^3 and R^4 when $n=0$, represents independently C1-4-alkyl or C3-7-cycloalkyl, optionally substituted by one or more halogen, hydroxy, lower alkyl and/or azido.

- 30 According to another preferred embodiment, the present invention relates to a compound as defined above wherein at least one of the R substituents chosen from the group R^2 , R^3 and R^4 when $n=1$ or from the group R^3 and R^4 when $n=0$, represents independently vinyl, optionally substituted by one or more halogen or/and lower alkyl.

- 35 According to another preferred embodiment, the present invention relates to a compound as defined above wherein at least one of the R substituents chosen from the group R^2 , R^3 and

R⁴ when n=1 or from the group R³ and R⁴ when n=0, represents independently ethynyl, propynyl or butynyl, optionally substituted by one or more halogen and/or lower alkyl.

According to another preferred embodiment, the present invention relates to a compound as defined above wherein at least one of the R substituents chosen from the group R², R³ and R⁴ when n=1 or from the group R³ and R⁴ when n=0, represents independently phenyl, optionally substituted by one or more halogen, lower alkyl, azido and/or nitro.

According to another preferred embodiment, the present invention relates to a compound as defined above wherein at least one of the R substituents chosen from the group R², R³ and R⁴ when n=1 or from the group R³ and R⁴ when n=0, represents independently 2- or 3-thienyl, optionally substituted by one or more halogen, acyl, cyano or/and lower alkyl.

According to a particular preferred embodiment, the present invention relates to a compound as defined above wherein at least one of the R substituents chosen from the group R³, R⁴ and R² when n=1 or from the group R³ and R⁴ when n=0, is hydroxymethyl, propyl, butyl, 3,3,3-trifluoropropyl, 2,2,2-trifluoroethyl, cyclopropylmethyl, iodomethyl, azidomethyl, 2-thienyl, 3-thienyl, phenyl, 3-chlorophenyl, 3-azidophenyl, 2,2-difluorovinyl, 2,2-dibromovinyl, 2,2-dichlorovinyl, 2-ethynyl, 5-methyl-2-thienyl, 5-formyl-2-ethynyl, 5-cyano-2-thienyl, 3-bromo-2-thienyl, 4-methyl-2-thienyl, 3,3,3-trifluoro-1-propynyl, 1-propynyl, cyclopropylethynyl, 3-methyl-1-butynyl, 1-butynyl, 2,2-difluoropropyl, 2-chloro-2,2-difluoroethyl, 2-bromo-2,2-difluoroethyl and 2-iodo-2,2-difluoroethyl.

According to yet another preferred embodiment, the present invention relates to a compound as defined above wherein R¹, R², R⁴ and R⁵ are hydrogen.

According to even another preferred embodiment, the present invention relates to a compound as defined above wherein R¹, R², R³ and R⁵ are hydrogen.

According to even another preferred embodiment, the present invention relates to a compound as defined above wherein n=1 and R¹, R³, R⁴ and R⁵ are hydrogen.

In all the above-mentioned scopes when the carbon atom to which R⁶ is attached is asymmetric it is preferably in the "S" - configuration.

Representative compounds of this invention as defined above are selected from the group consisting of 2-[5-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-5-propyl-1-piperidinyl)butanamide, 2-[2-oxo-5-(3,3,3-trifluoropropyl)-1-piperidinyl]butanamide, 2-[5-(cyclopropylmethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(iodomethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(azidomethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-5-phenyl-1-piperidinyl)butanamide, 2-[2-oxo-5-(2-thienyl)-1-piperidinyl]butanamide, 2-[2-oxo-5-(3-thienyl)-1-piperidinyl]butanamide, 2-[5-(3-chlorophenyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(3-azidophenyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2,2-difluorovinyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2,2-dibromovinyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2,2-

dichlorovinyl)-2-oxo-1-piperidinyl]butanamide, 2-(5-ethynyl-2-oxo-1-piperidinyl)butanamide, 2-[5-(5-methyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(5-formyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(5-cyano-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(3-bromo-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(4-methyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[2-oxo-5-(3,3,3-trifluoro-1-propynyl)-1-piperidinyl]butanamide, 2-[2-oxo-5-(1-propynyl)-1-piperidinyl]butanamide, 2-[5-(cyclopropylethynyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(3-methyl-1-butynyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(1-butynyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2,2-difluoropropyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2-chloro-2,2-difluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2-bromo-2,2-difluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-4-propyl-1-piperidinyl)butanamide, 2-[2-oxo-4-(3,3,3-trifluoropropyl)-1-piperidinyl]butanamide, 2-[4-(cyclopropylmethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(iodomethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(azidomethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-4-phenyl-1-piperidinyl)butanamide, 2-[2-oxo-4-(2-thienyl)-1-piperidinyl]butanamide, 2-[2-oxo-4-(3-thienyl)-1-piperidinyl]butanamide, 2-[4-(3-chlorophenyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(3-azidophenyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2-difluorovinyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2-dibromovinyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2-dichlorovinyl)-2-oxo-1-piperidinyl]butanamide, 2-(4-ethynyl-2-oxo-1-piperidinyl)butanamide, 2-[4-(5-methyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(5-formyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(5-cyano-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(3-bromo-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(4-methyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[2-oxo-4-(3,3,3-trifluoro-1-propynyl)-1-piperidinyl]butanamide, 2-[2-oxo-4-(1-propynyl)-1-piperidinyl]butanamide, 2-[4-(cyclopropylethynyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(3-methyl-1-butynyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(1-butynyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2-difluoropropyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2-chloro-2,2-difluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2-bromo-2,2-difluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2,2-trifluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-5-propyl-1-azepanyl)butanamide, 2-[2-oxo-5-(3,3,3-trifluoropropyl)-1-azepanyl]butanamide, 2-[5-(cyclopropylmethyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(iodomethyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(azidomethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-5-phenyl-1-azepanyl)butanamide, 2-[2-oxo-5-(2-thienyl)-1-azepanyl]butanamide, 2-[2-oxo-5-(3-thienyl)-1-azepanyl]butanamide, 2-[5-(3-chlorophenyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(3-azidophenyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2-difluorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2-dibromovinyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2-dichlorovinyl)-2-oxo-1-azepanyl]butanamide, 2-(5-ethynyl-2-oxo-

1-azepanyl]butanamide, 2-[5-(5-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(5-formyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(5-cyano-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(3-bromo-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(4-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[2-oxo-5-(3,3,3-trifluoro-1-propynyl)-1-azepanyl]butanamide, 2-[2-oxo-5-(1-propynyl)-1-azepanyl]butanamide, 2-[5-(cyclopropylethynyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(3-methyl-1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2-difluoropropyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2-chloro-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2-bromo-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2,2-trifluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-6-propyl-1-azepanyl]butanamide, 2-[2-oxo-6-(3,3,3-trifluoropropyl)-1-azepanyl]butanamide, 2-[6-(cyclopropylmethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(iodomethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(azidomethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-6-phenyl-1-azepanyl]butanamide, 2-[2-oxo-6-(2-thienyl)-1-azepanyl]butanamide, 2-[2-oxo-6-(3-thienyl)-1-azepanyl]butanamide, 2-[6-(3-chlorophenyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(3-azidophenyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2-difluorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2-dibromovinyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2-dichlorovinyl)-2-oxo-1-azepanyl]butanamide, 2-(6-ethynyl-2-oxo-1-azepanyl]butanamide, 2-[6-(5-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(5-formyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(5-cyano-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(3-bromo-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(4-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[2-oxo-6-(3,3,3-trifluoro-1-propynyl)-1-azepanyl]butanamide, 2-[2-oxo-6-(1-propynyl)-1-azepanyl]butanamide, 2-[6-(cyclopropylethynyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(3-methyl-1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2-difluoropropyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2-chloro-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2-bromo-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2,2-trifluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-4-propyl-1-azepanyl]butanamide, 2-[2-oxo-4-(3,3,3-trifluoropropyl)-1-azepanyl]butanamide, 2-[4-(cyclopropylmethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(iodomethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(azidomethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-4-phenyl-1-azepanyl]butanamide, 2-[2-oxo-4-(2-thienyl)-1-azepanyl]butanamide, 2-[2-oxo-4-(3-thienyl)-1-azepanyl]butanamide, 2-[4-(3-chlorophenyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(3-azidophenyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2-difluorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2-dibromovinyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2-dichlorovinyl)-2-oxo-1-azepanyl]butanamide, 2-(4-ethynyl-2-oxo-1-azepanyl]butanamide, 2-[4-(5-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(5-formyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(5-cyano-2-thienyl)-2-oxo-1-azepanyl]butanamide,

2-[4-(3-bromo-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(4-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[2-oxo-4-(3,3,3-trifluoro-1-propynyl)-1-azepanyl]butanamide, 2-[2-oxo-4-(1-propynyl)-1-azepanyl]butanamide, 2-[4-(cyclopropylethynyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(3-methyl-1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2-difluoropropyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2-chloro-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2-bromo-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2,2-trifluoroethyl)-2-oxo-1-azepanyl]butanamide.

The best results have been obtained with the following compounds :

- (2S)-2-[5-(iodomethyl)-2-oxo-1-piperidiny]butanamide,
- (2S)-2-[5-(azidomethyl)-2-oxo-1-piperidiny]butanamide,
- 2-(2-oxo-5-phenyl-1-piperidiny]butanamide,
- (2S)-2-[4-(iodomethyl)-2-oxo-1-piperidiny]butanamide,
- 2-[5-(iodomethyl)-2-oxo-1-azepanyl]butanamide.

It has now been found that compounds of formula I and their pharmaceutically acceptable salts are useful in a variety of pharmaceutical indications.

For example, the compounds according to the invention are useful for the treatment of epilepsy, epileptogenesis, seizure disorders and convulsions.

These compounds may also be used for the treatment of other neurological disorders including bipolar disorders, mania, depression, anxiety, migraine, trigeminal and other neuralgia, chronic pain, neuropathic pain, cerebral ischemia, cardiac arrhythmia, myotonia, cocaine abuse, stroke, myoclonus, essential tremor, dyskinesia and other movement disorders, neonatal cerebral haemorrhage, amyotrophic lateral sclerosis, spasticity, Parkinson's disease and other neurodegenerative diseases.

In addition the compounds according to the invention may be used in the treatment of bronchial asthma, asthmatic status and allergic bronchitis, asthmatic syndrome, bronchial hyperreactivity and bronchospastic syndromes as well as allergic and vasomotor rhinitis and rhinoconjunctivitis.

Thus, the present invention, in a further aspect, concerns the use of a compound of formula I or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of neurological and other disorders such as mentioned above.

In particular, the present invention concerns the use of a compound of formula I or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of epilepsy, bipolar disorders, chronic pain or neuropathic pain, migraine, bronchial-, asthmatic- or allergic conditions.

In particular, the present invention concerns the use of a compound as defined above for

the manufacture of a medicine for the treatment of epilepsy, epileptogenesis, seizure disorders and convulsions.

The activity and properties of the active compounds, oral availability and stability in vitro or in vivo can vary significantly among the optical isomers of the disclosed compounds.

5 In a preferred embodiment, the active compound is administered in an enantiomerically enriched form, i.e., substantially in the form of one isomer.

The present invention also concerns a method for treating epilepsy, epileptogenesis, seizure disorders, convulsions and other neurological disorders including bipolar disorders, mania, depression, anxiety, migraine, trigeminal and other neuralgia, chronic pain, neuropathic
10 pain, cerebral ischemia, cardiac arrhythmia, myotonia, cocaine abuse, stroke, myoclonus, essential tremor, dyskinesia and other movement disorders, neonatal cerebral haemorrhage, amyotrophic lateral sclerosis, spasticity, Parkinson's disease and other degenerative diseases, bronchial asthma, asthmatic status and allergic bronchitis, asthmatic syndrome, bronchial hyperreactivity and bronchospastic syndromes as well as allergic and vasomotor rhinitis and
15 rhinoconjunctivitis, in a mammal in need of such treatment, comprising administering a therapeutic dose of at least one compound as defined above.

The present invention also concerns a method for treating epilepsy, migraine, bipolar disorders, chronic pain or neuropathic pain or bronchial-, asthmatic- or allergic conditions, in a mammal in need of such treatment, comprising administering a therapeutic dose of at least one
20 compound of formula I or a pharmaceutically acceptable salt thereof to a patient.

The present invention also concerns a method for treating epilepsy, epileptogenesis, seizure disorders and convulsions.

The methods of the invention comprise administration to a mammal (preferably human) suffering from above mentioned conditions or disorders, of a compound according to the
25 invention in an amount sufficient to alleviate or prevent the disorder or condition.

The compound is conveniently administered in any suitable unit dosage form, including but not limited to one containing 5 to 1000 mg, preferably 25 to 500 mg of active ingredient per unit dosage form.

The term "treatment" as used herein includes curative treatment and prophylactic
30 treatment.

By "curative" is meant efficacy in treating a current symptomatic episode of a disorder or condition.

By "prophylactic" is meant prevention of the occurrence or recurrence of a disorder or condition.

35 The term "epilepsy" as used herein refers to a disorder of brain function characterised by the periodic and unpredictable occurrence of seizures. Seizures can be "non-epileptic" when

evoked in a normal brain by treatments such as electroshock or chemical convulsants or “epileptic” when evoked without evident provocation.

The term “seizure” as used herein refers to a transient alteration of behaviour due to the disordered, synchronous, and rhythmic firing of populations of brain neurones.

5 The term “migraine” as used herein means a disorder characterised by recurrent attacks of headache that vary widely in intensity, frequency, and duration. The attacks are commonly unilateral and are usually associated with anorexia, nausea, vomiting, phonophobia, and/or photophobia. In some cases they are preceded by, or associated with, neurological and mood disturbances. Migraine headache may last from 4 hours to about 72 hours. The International
10 Headache Society (IHS, 1988) classifies migraine with aura (classical migraine) and migraine without aura (common migraine) as the major types of migraine. Migraine with aura consists of a headache phase preceded by characteristic visual, sensory, speech, or motor symptoms. In the absence of such symptoms, the headache is called migraine without aura.

 The term “bipolar disorders” as used herein refers to those disorders classified as Mood
15 Disorders according to the Diagnostic and Statistical Manual of Mental Disorders, 4th edition (Diagnostic and Statistical Manual of Mental Disorders (DSM-IV TM), American Psychiatry Association, Washington, DC, 1994). Bipolar disorders are generally characterised by spontaneously triggered repeated (i.e. at least two) episodes in which the patient’s hyperexcitability, activity and mood are significantly disturbed, this disturbance consisting on
20 some occasions of an elevation of mood and increased energy and activity (mania or hypomania), and in other occasions a lowering of mood and decreased energy and activity (depression). Bipolar disorders are separated into four main categories in the DSM-IV (bipolar I disorder, bipolar II disorder, cyclothymia, and bipolar disorders not otherwise specified).

 The term “manic episode”, as used herein refers to a distinct period during which there is
25 an abnormally and persistently elevated, expansive, or irritable mood with signs of pressured speech and psychomotor agitation.

 The term “hypomania”, as used herein refers to a less extreme manic episode, with lower grade of severity.

 The term “major depressive episode”, as used herein refers to a period of at least 2 weeks
30 during which there is either depressed mood or the loss of interest or pleasure in nearly all activities with signs of impaired concentration and psychomotor retardation.

 The term “mixed episode”, as used herein refers to a period of time (lasting at least 1 week) in which the criteria are met both for a manic episode and for a major depressive episode nearly every day.

35 The term “chronic pain” as used herein refers to the condition gradually being recognised as a disease process distinct from acute pain. Conventionally defined as pain that persists

beyond the normal time of healing, pain can also be considered chronic at the point when the individual realises that the pain is going to be a persistent part of their lives for the foreseeable future. It is likely that a majority of chronic pain syndromes involves a neuropathic component, which is usually harder to treat than acute somatic pain.

5 The term "neuropathic pain" as used herein refers to pain initiated by a pathological change in a nerve which signals the presence of a noxious stimulus when no such recognisable stimulus exists, giving rise to a false sensation of pain. In other words, it appears that the pain system has been turned on and cannot turn itself off.

10 The activity of the compounds of formula I, or their pharmaceutically acceptable salts, as anticonvulsants can be determined in the audiogenic seizures model. The objective of this test is to evaluate the anticonvulsant potential of a compound by means of audiogenic seizures induced in sound-susceptible mice, a genetic animal model with reflex seizures. In this model of primary generalised epilepsy, seizures are evoked without electrical or chemical stimulation and the seizure types are, at least in part, similar in their clinical phenomenology to seizures occurring in
15 man (Löscher W. & Schmidt D., *Epilepsy Res.* (1998), 2, 145-181; Buchhalter J.R., *Epilepsia* (1993), 34, S31-S41). Results obtained with compounds of formula I are indicative of a strong pharmacological effect (Table III A, III B).

 Another assay indicative of potential anticonvulsant activity is binding to levetiracetam binding site (LBS) as hereinafter described (Table III A, III B).

20 The activity of the compounds of formula I, or their pharmaceutically acceptable salts, in chronic neuropathic pain can be determined in animal models. For example, chronic neuropathic pain can be modelled by pharmacologically inducing diabetes in rats. In this model, animals show progressive hyperalgesia to nociceptive stimuli, a symptom generally observed in patients with painful peripheral neuropathy (Courteix C, Eschalier, A. and Lavarenne J., *Pain*
25 (1993), 53, 81-88). This model was shown to possess a high pharmacological predictivity (Courteix C, Bardin M., Chantelauze C., Lavarenne J and Eschalier, A., *Pain*, 57 (1994) 153-160).

 The activity of the compounds of formula I, or their pharmaceutically acceptable salts, in bipolar disorders can be assessed in animal models. For example, bipolar disorders and
30 especially mania can be modelled by pharmacologically inducing hyperactivity in rats and evaluating their behaviour in an Y maze. In such a situation, therapeutic agents effective in man, like Lithium and sodium valproate decrease the hyperactivity, thus validating the predictivity of the model (Cao B. J. and Peng N. A., *Eur. J. Pharmacol.* (1993), 237, 177-181; Vale A. L. and Ratcliffe F. *Psychopharmacology*, 91 (1987) 352-355).

35 Potential anti-asthmatic properties of the compounds of formula I, or their pharmaceutically acceptable salts would be tested for in an animal model of allergic asthma, in

which guinea pigs sensitised to ovalbumin are challenged with the antigen and investigated for changes in pulmonary function and airway inflammatory cell content. (Yamada et al. (1992) Development of an animal model of late asthmatic response in guinea pigs and effects anti-asthmatic drugs. Prostaglandins, 43: 507-521).

5 Activity in any of the above mentioned indications can of course be determined by carrying out suitable clinical trials in a manner known to a person skilled in the relevant art for the particular indication and/or in the design of clinical trials in general.

For treating diseases, compounds of formula I or their pharmaceutically acceptable salts may be employed at an effective daily dosage and administered in the form of a pharmaceutical
10 composition.

Therefore, another embodiment of the present invention concerns a pharmaceutical composition comprising an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof in combination with a pharmaceutically acceptable diluent or carrier.

To prepare a pharmaceutical composition according to the invention, one or more of the
15 compounds of formula I or a pharmaceutically acceptable salt thereof, is intimately admixed with a pharmaceutical diluent or carrier according to conventional pharmaceutical compounding techniques known to the skilled practitioner.

Suitable diluents and carriers may take a wide variety of forms depending on the desired route of administration, e.g., oral, rectal, or parenteral.

20 Pharmaceutical compositions comprising compounds according to the invention can, for example, be administered orally or parenterally, i.e., intravenously, intramuscularly or subcutaneously, intrathecally.

Pharmaceutical compositions suitable for oral administration can be solids or liquids and can, for example, be in the form of tablets, pills, dragees, gelatine capsules, solutions, syrups,
25 and the like.

To this end the active ingredient may be mixed with an inert diluent or a non-toxic pharmaceutically acceptable carrier such as starch or lactose. Optionally, these pharmaceutical compositions can also contain a binder such as microcrystalline cellulose, gum tragacanth or gelatine, a disintegrant such as alginic acid, a lubricant such as magnesium stearate, a glidant
30 such as colloidal silicon dioxide, a sweetener such as sucrose or saccharin, or colouring agents or a flavouring agent such as peppermint or methyl salicylate.

The invention also contemplates compositions which can release the active substance in a controlled manner. Pharmaceutical compositions which can be used for parenteral administration are in conventional form such as aqueous or oily solutions or suspensions
35 generally contained in ampoules, disposable syringes, glass or plastics vials or infusion containers.

In addition to the active ingredient, these solutions or suspensions can optionally also contain a sterile diluent such as water for injection, a physiological saline solution, oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents, antibacterial agents such as benzyl alcohol, antioxidants such as ascorbic acid or sodium bisulphite, chelating agents such as ethylene diamine-tetra-acetic acid, buffers such as acetates, citrates or phosphates and agents for adjusting the osmolarity, such as sodium chloride or dextrose.

These pharmaceutical forms are prepared using methods which are routinely used by pharmacists.

The amount of active ingredient in the pharmaceutical compositions can fall within a wide range of concentrations and depends on a variety of factors such as the patient's sex, age, weight and medical condition, as well as on the method of administration. Thus the quantity of compound of formula I in compositions for oral administration is at least 0.5 % by weight and can be up to 80 % by weight with respect to the total weight of the composition.

In accordance with the invention it has also been found that the compounds of formula I or the pharmaceutically acceptable salts thereof can be administered alone or in combination with other pharmaceutically active ingredients. Non-limiting examples of such additional compounds which can be cited for use in combination with the compounds according to the invention are antivirals, antispastics (e.g. baclofen), antiemetics, antimanic mood stabilizing agents, analgesics (e.g. aspirin, ibuprofen, paracetamol), narcotic analgesics, topical anaesthetics, opioid analgesics, lithium salts, antidepressants (e.g. mianserin, fluoxetine, trazodone), tricyclic antidepressants (e.g. imipramine, desipramine), anticonvulsants (e.g. valproic acid, carbamazepine, phenytoin), antipsychotics (e.g. risperidone, haloperidol), neuroleptics, benzodiazepines (e.g. diazepam, clonazepam), phenothiazines (e.g. chlorpromazine), calcium channel blockers, amphetamine, clonidine, lidocaine, mexiletine, capsaicin, caffeine, quetiapine, serotonin antagonists, β -blockers, antiarrhythmics, triptans, ergot derivatives.

Of particular interest in accordance with the present invention are combinations of at least one compound of formula I or a pharmaceutically acceptable salt thereof and at least one compound inducing neural inhibition mediated by GABA_A receptors. The compounds of formula I exhibit a potentiating effect on the compounds inducing neural inhibition mediated by GABA_A receptors enabling, in many cases, effective treatment of conditions and disorders under reduced risk of adverse effects.

Examples of compounds inducing neural inhibition mediated by GABA_A receptors include the following: benzodiazepines, barbiturates, steroids, and anticonvulsants such as valproate, vigabatrin, tiagabine or pharmaceutical acceptable salts thereof.

Benzodiazepines include the 1,4 benzodiazepines, such as diazepam and clonazepam, and the 1,5-benzodiazepines, such as clobazam. Preferred compound is clonazepam.

Barbiturates include phenobarbital and pentobarbital. Preferred compound is phenobarbital.

Steroids include adrenocorticotrophic hormones such as tetracosactide acetate, etc.

Anticonvulsants include hydantoins (phenytoin, ethotoin, etc), oxazolidines (trimethadione, etc.), succinimides (ethosuximide, etc.), phenacemides (phenacemide, acetylpheneturide, etc.), sulfonamides (sulthiame, acetazolamide, etc.), aminobutyric acids (e.g. gamma-amino-beta-hydroxybutyric acid, etc.), sodium valproate and derivatives, carbamazepine and so on.

Preferred compounds include valproic acid, valpromide, valproate pivoxil, sodium valproate, semi-sodium valproate, divalproex, clonazepam, phenobarbital, vigabatrin and tiagabine.

For the preferred oral compositions, the daily dosage is in the range 5 to 1000 milligrams (mg) of compounds of formula I.

In compositions for parenteral administration, the quantity of compound of formula I present is at least 0.5 % by weight and can be up to 33 % by weight with respect to the total weight of the composition. For the preferred parenteral compositions, the dosage unit is in the range 5 to 1000 mg of compounds of formula I.

The daily dose can fall within a wide range of dosage units of compound of formula I and is generally in the range 5 to 1000 mg. However, it should be understood that the specific doses can be adapted to particular cases depending on the individual requirements, at the physician's discretion.

The amount of the active ingredients (compound I and compound inducing neural inhibition mediated by the GABA_A receptors) in the pharmaceutical composition of the invention will vary depending on the mammal to which the compositions are administered, the disease to be treated, other active ingredients present, etc. Generally, the amount of the compound inducing neural inhibition mediated by the GABA_A receptors and the amount of compound I for a given composition and dosage form can be readily determined employing routine procedures.


According to another aspect, the invention relates to several processes for preparing a compound having the general formula (I) as described above.

The following process description sets forth certain synthesis routes in an illustrative manner. Other alternative and/or analogous methods will be readily apparent to those skilled in this art. As used herein in connection with substituent meanings, "=" means "is" and "≠" means "is other than".

According to one embodiment, compounds having the general formula (I), wherein A¹ = O and X = -CONR⁷R⁸ or -COOR⁹, may be made by the cyclisation of an aminoester of formula (A) wherein Q¹ represents a linear or branched C1-4-alkyl. Said aminoester of formula (A)

wherein $n = 0$ and $R^3 = \text{COOR}^{13}$ or wherein $n = 1$ and $R^2 = \text{COOR}^{13}$, wherein R^{13} represents a linear or branched alkyl group optionally optically active, are obtained by reaction of a compound of formula (B) with an unsaturated diester derivative of formula (C) or (D), respectively, according to the following scheme 1.

5




Scheme 1

The first reaction of compound (B) with compound (C) or (D) leading to the formation of the compound of formula (A) is described in: Street, L.J., Baker, R., Book, T., Kneen, C.O., MacLeod, A.M., Merchant, K.J., Showell, G. A., Saunders, J., Herbert, R. H., Freedman, S.B., Harley, E.A., J. Med. Chem. (1990), 33, 2690-2697.

When Q^1 represents a methyl or ethyl, the second reaction converting compound of formula (A) into (I) is known by the man skilled in the art and is generally carried out between room temperature and 150°C , in the presence or not of a catalyst such as acetic acid, hydroxybenzotriazole or 2-hydroxypyridine. When Q^1 does not represent a methyl or ethyl, an ester of formula (A) is hydrolysed under acidic or basic conditions and then cyclised under conventional peptide synthesis conditions, by using coupling agents, for example dicyclohexylcarbodiimide (Bodanszky, M., Bodanszky, A., in "The Practice of Peptide Synthesis", Springer Verlag, 1984).

According to another embodiment, compounds of the general formula (I), wherein $\text{A}^1 = \text{O}$ and $\text{X} = -\text{CONR}^7\text{R}^8$ and $n = 0$ may conveniently be made by reaction of a compound of formula

(E), wherein X^1 represents a halogen atom, preferably an iodine or a chlorine atom, and X^2 represents $-OQ^1$ or a halogen atom, preferably a chlorine atom, with a compound of formula (F). Said compound of formula (E) may be obtained by opening of a lactone of formula (G) in the presence of a halogenating agent, for instance TMSI or $SOCl_2/ZnCl_2$, followed if necessary by halogenation of the obtained halogenoacid ($X^2 = OH$). Furthermore, said compound of formula (G) may be prepared by reaction of a compound of formula (J) with an organolithium compound of formula R^4Li in the presence of CuI or, when in formula (G) $R^2 = H$, by reduction of a carboxylate of formula (H) wherein Q^4 is methyl or ethyl, in the presence of a borohydride reagent. All these reactions are depicted schematically in the following scheme 2.



Scheme 2


The reaction of the compound of formula (J) with the organolithium compound of formula R^4Li in the presence of CuI may be carried out according to the procedures described in: Alexakis, A., Berlan, J., Besace, Y., Tetrahedron Lett. (1986), 27, 1047-1050 and in: Lipshutz, B.H., Ellsworth, E.L., Siahaan, T., J. Am. Chem. Soc. (1989), 111, 1351-1358.

Opening of the lactone of formula (G) may be performed according to the procedure described in: Mazzini, C., Lebreton, J., Alphand, V., Furstoss, R., Tetrahedron Lett. (1998), 38, 1195-1196 and in Olah, G.A., Narang, S.C., Gupta, B.G.B., Malhotra, R., J. Org. Chem. (1979), 44, 1247-1250. Halogenation ($X^2 = \text{halogen}$) or esterification ($X^2 = OQ^1$) of the obtained haloalkanoic acid ($X^2 = OH$) may be performed under any conditions known to the person skilled in the art.

The reaction of a compound of formula (E) with a compound of formula (F) forming compound (I) may be carried out in an inert solvent, for example dichloromethane, at a

temperature of about 0 °C in the presence of a phase transfer catalyst, such as tetrabutylammonium bromide in the presence of powdered potassium hydroxide (Patent Application GB 2225322 A).

According to yet another embodiment, compounds of the general formula (I), wherein $A^1 = O$ and $X = -COOR^9$, R^9 representing an alkyl group, may also be prepared by reaction of a compound of formula (K) wherein M represents an alkali metal, with a compound of formula (L) wherein X^3 represents a halogen atom, according to the following scheme 3.



Scheme 3

Compounds of formula (K) may be prepared according to the procedure described in: Georg, G., Guan, X., Kant, J., Bioorg. Med. Chem. Lett. (1991), 1, 125.

Alternatively, compounds of formula (K) wherein $n = 0$ may be prepared according to the procedure described in: Koelsch, C.F., J. Am. Chem. Soc. (1943), 65, 2093-2095.

According to yet a further embodiment, the invention relates to a process for preparing a compound of general formula (I), wherein $A^1 = O$ and $X = -CONR^7R^8$ and none of the groups R^1, R^2, R^3, R^4, R^5 and R^6 being substituted by carboxylic or sulfonic acids or esters or ketone, wherein said process comprises the ammonolysis of the corresponding compound wherein $X = -COOR^9$, R^9 being a hydrogen atom or a linear or branched C1-4-alkyl group. Alternatively, transformation into the amino form may be done under conventional peptidic synthesis conditions using coupling agents, for example alkyl chloroformate or dicyclohexylcarbodiimide.

Other processes for preparation of compounds of the general formula (I) and intermediates are known by the man skilled in the art.

For instance, compounds of formula (I) wherein $A^1 = O$, $X = CONR^7R^8$, and one of the groups R^1, R^2, R^3, R^4, R^5 and R^6 represents $-G^3-COOCH_3$, G^3 being a bond or an alkylene group, are key synthesis intermediates for corresponding compounds wherein one of the groups R^1, R^2, R^3, R^4, R^5 and R^6 represents $-G^3-CH_2OH$ (March, J., "Advanced Organic Chemistry, Third Edition", John Wiley & Sons, (1985), 1101-1102).

Furthermore, compounds of formula (I) wherein $A^1 = O$, $X = \text{CONR}^7\text{R}^8$, and one of the groups R^1 , R^2 , R^3 , R^4 , R^5 and R^6 represents $-\text{G}^3-\text{CH}_2\text{OH}$, G^3 being a bond or an alkylene group, are key synthesis intermediates for corresponding compounds wherein one of the groups R^1 , R^2 , R^3 , R^4 , R^5 and R^6 represents $-\text{G}^3-\text{CH}_2\text{X}^3$ or $-\text{G}^3-\text{CHO}$, wherein X^3 represents a
 5 chlorine, a bromine or a iodine atom, or a group of formula $-\text{O}-\text{SO}_2-\text{R}^{14}$ wherein R^{14} represents an alkyl or an aryl group. These transformations may be performed under any conditions known to the person skilled in the art.

In addition, compounds of formula I wherein $A^1 = O$, $X = \text{CONR}^7\text{R}^8$, and one of the groups R^1 , R^2 , R^3 , R^4 , R^5 and R^6 represents $-\text{G}^3-\text{CH}_2\text{OSO}_2\text{R}^{14}$, G^3 being a bond or an
 10 alkylene group and R^{14} being a methyl or a tolyl group, are key synthesis intermediates for corresponding compounds wherein one of the groups R^1 , R^2 , R^3 , R^4 , R^5 and R^6 represents $-\text{G}^3-\text{CH}_2\text{N}_3$ or $-\text{G}^3-\text{CH}_2\text{X}^4$, wherein X^4 represents an halogen atom. These transformations may be performed under any conditions known to the person skilled in the art.

Finally, compounds of formula I wherein $A^1 = O$, $X = \text{CONR}^7\text{R}^8$, and one of the groups
 15 R^1 , R^2 , R^3 , R^4 , R^5 and R^6 represents $-\text{G}^3-\text{CHO}$, G^3 being a bond or an alkylene group, are key synthesis intermediates for corresponding compounds wherein one of the groups R^1 , R^2 , R^3 , R^4 , R^5 and R^6 represents $-\text{G}^3-\text{R}^{15}$ wherein R^{15} represents a vinyl group not substituted, mono- or di-substituted by halogen atoms or alkyl groups (Wittig type reaction).

According to a last embodiment, the present invention encompasses any of the
 20 intermediates prepared by any of the processes described above.

Representative intermediates of this invention as defined above are selected from the group consisting of methyl 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-(3S)-3-piperidinecarboxylate and methyl 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-(3R)-3-piperidinecarboxylate and their mixtures, (2S)-2-[(5R)-5-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide and (2S)-2-[(5S)-5-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide and their mixtures, 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-(3R)-3-piperidinyl)methyl methanesulfonate and {1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-(3S)-3-piperidinyl)methyl methanesulfonate and their mixtures, 5-phenyl-2-piperidinone, ethyl 2-(2-oxo-5-phenyl-1-piperidinyl)butanoate, ethyl 7-oxo-4-azepanecarboxylate, ethyl 1-[1-(tert-butoxycarbonyl)propyl]-7-oxo-4-azepanecarboxylate, 2-[5-(ethoxycarbonyl)-2-oxo-1-azepanyl]butanoic acid, 1-[1-(aminocarbonyl)propyl]-7-oxo-4-azepanecarboxylate, 2-[5-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide.
 25
 30

In the preparation processes according to the invention, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example extraction, crystallization, distillation,
 35 trituration and chromatography, or any combination of the same.

When compounds of formula I present one or several stereogenic centres, and that non-stereoselective methods of synthesis are used, resolution of the mixture of stereoisomers can best be effected in one or several steps, involving generally sequential separation of mixtures of diastereomers into their constituting racemates, using preferably chromatographic separations on achiral or chiral phase in reversed or preferably in direct mode, followed by at least one ultimate step of resolution of each racemate into its enantiomers, using most preferably chromatographic separation on chiral phase in reversed or preferably in direct mode. Alternatively, when partly stereoselective methods of synthesis are used, the ultimate step may be a separation of diastereomers using preferably chromatographic separations on achiral or chiral phase in reversed or preferably in direct mode.

Some of the intermediates are known compounds or may be prepared according to art-known procedures.

The present compounds of this invention differ from the art by their structure and by their favourable pharmacological properties. The following examples are provided for illustrative purposes only and are not intended, nor should they be construed, as limiting the invention in any manner. Those skilled in the art will appreciate that routine variations and modifications of the following examples can be made without exceeding the spirit or scope of the invention.

EXAMPLES

EXAMPLE 1 : Analytical characterization of 6- and 7-ring compounds

Unless specified otherwise in the examples, characterization of the compounds was performed according to the following methods:

NMR spectra are recorded on a BRUKER AC 250 Fourier Transform NMR Spectrometer fitted with an Aspect 3000 computer and a 5 mm $^1\text{H}/^{13}\text{C}$ dual probehead or BRUKER DRX 400 FT NMR fitted with a SG Indigo² computer and a 5 mm inverse geometry $^1\text{H}/^{13}\text{C}/^{15}\text{N}$ triple probehead. The compound is studied in DMSO- d_6 (or CDCl_3) solution at a probe temperature of 313 K and at concentrations ranging from 2 to 20 mg/ml. The instrument is locked on the deuterium signal of DMSO- d_6 (or CDCl_3). Chemical shifts are given in ppm downfield from TMS taken as internal standard.

Mass spectrometric measurements in LC/MS mode are performed as follows :

HPLC conditions

Analyses are performed using a WATERS Alliance HPLC system mounted with an INERTSIL ODS 3-, DP 5 μm , 250 X 4.6 mm column.

The gradient runs from 100 % solvent A (acetonitrile, water, TFA (10/90/0.1, v/v/v)) to 100 % solvent B (acetonitrile, water, TFA (90/10/0.1, v/v/v)) in 7 min with a hold at 100 % B of 4 min. The flow rate is set at 2.5 ml/min and a split of 1/10 is used just before API source. The chromatography is carried out at 30 °C.

MS conditions

Samples are dissolved in acetonitrile/water, 70/30, v/v at the concentration of about 250 μ g/ml. API spectra (+ or -) are performed using a FINNIGAN (San Jose, CA, USA) LCQ ion trap mass spectrometer. APCI source operates at 450 °C and the capillary heater at 160 °C. ESI
5 source operates at 3.5 kV and the capillary heater at 210 °C.

Mass spectrometric measurements in EI/DIP mode are performed as follows: samples are vaporized by heating the probe from 50 °C to 250 °C in 5 min. EI (Electron Impact) spectra are recorded using a FINNIGAN (San Jose, CA, USA) TSQ 700 tandem quadrupole mass spectrometer. The source temperature is set at 150 °C.

10 Specific rotation is recorded on a Perkin-Elmer MC241 or MC341 polarimeter. The angle of rotation is recorded at 25 °C on 1 % solutions in MeOH. For some molecules, the solvent is CH_2Cl_2 or DMSO, due to solubility problems.

Water content is determined using a Metrohm microcoulometric Karl Fischer titrator.

Preparative chromatographic separations are performed on silicagel 60 Merck, particle
15 size 15-40 μ m, reference 1.15111.9025, using in-house modified Jobin Yvon-type axial compression columns (80 mm i.d.), flow rates between 70 and 150 ml/min. Amount of silicagel and solvent mixtures are as described in individual procedures.


Preparative chiral chromatographic separations are performed on a DAICEL Chiralpak AD 20 μ m, 100*500 mm column using an in-house build instrument with various mixtures of
20 lower alcohols and C_5 to C_8 linear, branched or cyclic alkanes at \pm 350 ml/min. Solvent mixtures are as described in individual procedures.

Melting points are determined on a Büchi 535 Totoli-type fusionometre, and are not corrected, or by the onset temperature on a Perkin Elmer DSC 7.

Unless specified otherwise in the examples, the compounds are obtained in the neutral
25 form.

EXAMPLE 2 : Preparation of intermediates


2.1. Synthesis of methyl 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidinecarboxylate 1 and 2.



30 In a 100 ml three necked flask fitted with magnetic stirrer and reflux condenser, under inert atmosphere, 19 g (186 mmoles, 2.66 eq) of (S)-2-aminobutanamide 22 and 12 g (70

mmoles, 1 eq) of dimethyl 2-methylenepentanedioate 21 were dissolved in 50 ml of MeOH. 5 ml of acetic acid were added and the mixture was brought to reflux for 7 days. After cooling down, it was concentrated to dryness, dissolved in CH_2Cl_2 , washed successively with water, 0.5 N HCl and brine, dried over MgSO_4 and concentrated to dryness to give 17.8 g of crude amide ester as a mixture of diastereomers. It was purified by PrepLC (1 kg SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$, 97.25/2.5/0.25) to give 6.8 g of diastereoisomer 1 (first eluted) and 4.59 g of diastereoisomer 2 (second eluted), 67 % yield. Both were recrystallised from ethyl acetate/hexane.

2.2. Synthesis of (2S)-2-[5-(hydroxymethyl)-2-oxo-1-piperidiny]butanamide 3.




In a 250 ml three necked flask fitted with magnetic stirrer and reflux condenser, under inert atmosphere, 6.29 g (26 mmoles, 1 eq) of methyl 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidinecarboxylate 1 were dissolved in 35 ml of EtOH and the mixture cooled down to 0 °C. 2.5 g (66 mmoles, 9.5 eq) of solid NaBH_4 were then added by portions over 1.5 hour, all the while maintaining the temperature around 0 °C. After 2 hours, the temperature was raised to 12 °C for 1 hour, and lowered again to 0 °C. 10 ml of a saturated solution of NH_4Cl were added slowly, followed by 10 ml of acetone, and the mixture was left 1 hour at 0 °C. The mixture was filtered, the precipitate washed with 3 x 10 ml of EtOH and the combined organic fractions concentrated to dryness to give 5.9 g of crude alcohol. It was purified by PrepLC (500 g SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 90/10) to give 2.6 g of (2S)-2-[5-(hydroxymethyl)-2-oxo-1-piperidiny]butanamide 3, (52% yield).

Compound such as (2S)-2-[4-(hydroxymethyl)-2-oxo-1-piperidiny]butanamide can be synthesized in an analogous way.

2.3. Synthesis of 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidiny]methyl methanesulfonate 23.


30



In a 100 ml three necked flask fitted with magnetic stirrer, dropping funnel and reflux condenser under inert atmosphere, 1.9 g (8.8 mmol, 1 eq) of (2S)-2-[5-(hydroxymethyl)-2-oxo-1-piperidiny]butanamide **3** were suspended in 50 ml of acetone and cooled down to 0 °C. 2.5 ml (17.8 mmol, 2 eq) of dry triethylamine were added in one portion, followed by dropwise addition of a solution of 0.75 ml (8.8 mmol, 1 eq) of methanesulfonyl chloride in 5 ml of acetone, all the while maintaining the temperature below 4 °C. After 3 hours, the mixture was filtered, the residue washed with acetone and the combined organic phases concentrated to dryness to give 3.4 g of crude 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidiny]methyl methanesulfonate. It was purified by PrepLC (400 g SiO₂, CH₂Cl₂/MeOH, 96/4) to give 1.4 g of pure methanesulfonate **23**, (62 % yield).


Compound such as {1-[(1S)-1-(aminocarbonyl)propyl]-2-oxo-4-piperidiny]methyl methanesulfonate **24** can be synthesized in an analogous way.

EXAMPLE 3: synthesis of (2S)-2-[5-(azidomethyl)-2-oxo-1-piperidiny]butanamide **7**.




In a 50 ml three necked flask fitted with magnetic stirrer and reflux condenser, under inert atmosphere, 1.4 g (5.1 mmol, 1 eq) of 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidiny]methyl methanesulfonate **23** were dissolved in 15 ml of DMF. 400 mg (6.15 mmol, 1.2 eq) of solid sodium azide were added in one portion, and the mixture was heated up to 80 °C for 1 hour. The DMF was evaporated in vacuum, the residue suspended in CH₂Cl₂ and the precipitate filtered off. The filtrate was concentrated to dryness to give 1 g of crude material. It was purified by PrepLC (200 g SiO₂, CH₂Cl₂/MeOH, 97/3) to give 500 mg of pure azide **7**, 45 % yield.

EXAMPLE 4: synthesis of (2S)-2-[4-(iodomethyl)-2-oxo-1-piperidiny]butanamide **13**.



In a 150 ml three necked flask fitted with magnetic stirrer and reflux condenser, under inert atmosphere, 6.88 g (23.5 mmoles) of {1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidiny]methyl methanesulfonate 24 were dissolved in 86 ml of THF. 3.9 g (25.9 mmoles, 1.1 eq) of sodium iodide were added in one portion and the mixture was brought to reflux. After 5 hours, 780 mg (5.2 mmoles, 0.22 eq) of sodium iodide were added and reflux continued for a total of 10 hours. After cooling down to 0 °C, the mixture was filtered, the precipitate washed with THF, the combined organic fractions concentrated to dryness and purified by PrepLC (800 g SiO₂, CH₂Cl₂/MeOH, 95/5) to give 7.21 g of crude compound. It was crystallised from 80 ml ethanol to give 5 g of pure iodide 13, 66 % yield.

EXAMPLE 5: preparation of ethyl 2-(2-oxo-5-phenyl-1-piperidiny)butanoate 26.




Synthesis of the 5-phenyl-2-piperidinone 25: Koelsch, J. Amer. Chem. Soc. (1943), 65, 2093; the racemate was resolved by Chiral PrepLC (Chiralpak AD, petroleum ether/isopropanol, 90/10, 20 °C).

In a 1 l three necked flask fitted with reflux condenser, magnetic stirrer and dropping funnel under inert atmosphere, 9.8 g (87 mmoles) of potassium t-butoxide were suspended in 600 ml of dry toluene. 11.8 g (67 mmoles) of 5-phenyl-2-piperidinone 25 were added, and the mixture stirred for 30 min. 17 g (87 mmoles) of ethyl 2-bromobutanoate were added and the temperature raised to 70 °C for 5 hours. 600 ml of ethyl acetate were added, the solution was washed twice with brine, the organic phase dried over magnesium sulfate, filtered and concentrated to dryness. The residue was purified by PrepLC (1 kg SiO₂, CH₂Cl₂/EtOH, 97.5/2.5) to give 3.7 g of pure ester 26 as a mixture of diastereomers, 70.4 % yield.

Compound such as methyl 1-[1-(aminocarbonyl)propyl]-2-oxo-4-piperidinecarboxylate can be synthesized in an analogous way.


EXAMPLE 6: preparation of 2-(2-oxo-5-phenyl-1-piperidiny)butanamide 9, 10, 11 and

5 12.



In a 1 l three necked flask fitted with reflux condenser, magnetic stirrer and a gas inlet
10 tube dipping in the solution, 13.6 g (47 mmoles) of ethyl 2-(2-oxo-5-phenyl-1-
piperidiny)butanoate 26 were dissolved in 800 ml of methanol containing 22 mg of dissolved
sodium. Gaseous ammonia was bubbled through the solution, and the saturated solution kept
at room temperature for 3 days, while occasionally resaturating with ammonia. After completion
of the reaction, the solution was concentrated to dryness. The residue was purified by PrepLC (1
15 kg SiO₂, CH₂Cl₂/MeOH, 95/5) to give 10.1 g of butanamide. The diastereoisomers were then
separated (Chiralpak AD, ethanol) to give 4.5 g of diastereoisomer 11 (first eluted) and 4.3 g of
diastereoisomer 12 (second eluted) (72 %). Both were recrystallised from ethanol.

EXAMPLE 7: method for preparation of 7-ring compounds



7.1. Preparation of ethyl 7-oxo-4-azepanecarboxylate 28.

In a 250 ml three necked flask fitted with reflux condenser, magnetic stirrer and dropping funnel under inert atmosphere, 5.6 ml (5.98 g, 34 mmole, 1 eq) of ethyl 4-oxocyclohexanecarboxylate 27 were dissolved in 50 ml of CHCl_3 . 6.65 g (102 mmole, 3 eq) of sodium azide were added, followed by 22.1 ml (32.8 g, 341 mmole, 10 eq) of methanesulfonic acid dissolved in 20 ml of over 30 min. The mixture was heated to reflux for 1 hour and cooled down to 10 °C. 200 ml of saturated sodium bicarbonate solution were added, the mixture decanted and the aqueous phase extracted with 3 x 150 ml of CHCl_3 , the combined organic phases were dried over magnesium sulfate, filtered and concentrated to dryness. The residue was purified by PrepLC (800 g SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 97/3) to give 5.1 g of crude compound. It was crystallised from 40 ml of diethylether to give 4.41 g of pure ester 28, 70 % yield.

7.2. Preparation of ethyl 1-[1-(tert-butoxycarbonyl)propyl]-7-oxo-4-azepanecarboxylate 29.

In a 150 ml three necked flask fitted with reflux condenser, magnetic stirrer and dropping funnel under inert atmosphere, 5.09 g (27.5 mmole, 1 eq) of ethyl 7-oxo-4-azepanecarboxylate 28 and 12.27 g (55 mmole, 2 eq) of tert-butyl 2-bromobutanoate were dissolved in 70 ml of acetonitrile. The temperature was raised to 50 °C and 1.32 g (55 mmole, 2 eq) of sodium hydride were added portionwise (exothermic!). The mixture was stirred one more hour at 50 °C and concentrated to dryness, the residue was poured on ice/water and neutralised with solid

ammonium chloride. The mixture was extracted three times with CH_2Cl_2 , the combined organic phases were dried over magnesium sulfate, filtered and concentrated to dryness. The residue was purified by PrepLC (1 kg SiO_2 , CH_2Cl_2 /MTBE, 85/15) to give 8.5 g of pure diester 29, 69 % yield.

5

7.3. Preparation of 2-[5-(ethoxycarbonyl)-2-oxo-1-azepanyl]butanoic acid 30.

In a 25 ml three necked flask fitted with magnetic stirrer and dropping funnel under inert atmosphere, 1 g (3.1 mmoles) of pure ethyl 1-[1-(tert-butoxycarbonyl)propyl]-7-oxo-4-azepanecarboxylate 29 was dissolved in 5 ml of CH_2Cl_2 . The mixture was cooled down to 0 °C, 10 5 ml of trifluoroacetic acid was added. The mixture was kept at 0 °C under stirring for 24 hours, concentrated to dryness and used as such in the next step.

7.4. Preparation of ethyl 1-[1-(aminocarbonyl)propyl]-7-oxo-4-azepanecarboxylate 31.

In a 250 ml three necked flask fitted with reflux condenser, magnetic stirrer and 15 dropping funnel under inert atmosphere, 5.48 g (20.2 mmoles, 1 eq) of 2-[5-(ethoxycarbonyl)-2-oxo-1-azepanyl]butanoic acid 30 were dissolved in 150 ml of ethyl acetate, and 7.44 g (40.4 mmoles, 2 eq) of pentafluorophenol and 8.34 g (40.4 mmoles, 2 eq) of dicyclohexycarbodiimide were added, and the mixture kept under stirring at room temperature for 4 hours. 1.86 g of pentafluorophenol and 2.08 g of dicyclohexycarbodiimide (10.7 mmoles, 0.5 eq each) were 20 added, and stirring continued for 1 hour. The mixture was filtered, the residue washed with ethyl acetate, and the combined organic phases concentrated to dryness. The residue was taken up in 160 ml of CH_2Cl_2 , cooled down to 0 °C, saturated with gaseous ammonia and stirred at that temperature for 6 hours with occasional resaturations. The mixture was filtered, the residue washed with CH_2Cl_2 , and the combined organic phases concentrated to dryness. The residue 25 was purified and separated into diastereomers by PrepLC (1 kg SiO_2 , CH_2Cl_2 /i-PrOH/ NH_4OH , 92.5/7/0.5), to give 2.33 g of the first eluted diastereoisomer (racemate) and 2.7 g of the second diastereomer (racemate), 92 % total yield.

Steps 7.5. and 7.6. were described in the previous examples (2.2, 2.3 and 4).

30 Compounds with other ring substituents than iodine, or with substituents at other positions on the ring structure can be synthesized in an analogous way.

EXAMPLE 8 : Characterization of the interactions between a test substance and LBS.

LBS stands for Levetiracetam Binding Site, it is an unknown component of brain tissue 35 that has been shown to bind levetiracetam (M. Noyer et al., Eur. J. Pharmacol. (1995), 286, 137-146).

The inhibition constant (K_i) of a compound is determined in competitive binding experiments by measuring the binding of a single concentration of a radioactive ligand at equilibrium with various concentrations of the unlabeled test substance. The concentration of the test substance inhibiting 50 % of the specific binding of the radioligand is called the IC_{50} .

5 The equilibrium dissociation constant K_i is proportional to the IC_{50} and is calculated using the equation of Cheng and Prusoff (Cheng Y. et al., Biochem. Pharmacol. (1972), 22, 3099-3108).

The concentration range usually encompasses 6 log units with variable steps (0.3 to 1 log). Assays are performed in mono- or duplicate, each K_i determination is performed on two different samples of test substance. Cerebral cortex from 200-250 g male Sprague-Dawley rats
10 were homogenised using a Potter S homogeniser (10 strokes at 1,000 rpm; Braun, Germany) in 20 mmol/l Tris-HCl (pH 7.4), 250 mmol/l sucrose (buffer A); all operations were performed at 4 °C. The homogenate was centrifuged at 30,000 g for 15 min. The crude membrane pellet obtained was resuspended in 50 mmol/l Tris-HCl (pH 7.4), (buffer B) and incubated 15 min at 37 °C, centrifuged at 30,000 g for 15 min and washed twice with the same buffer. The final pellet
15 was resuspended in buffer A at a protein concentration ranging from 15 to 25 mg/ml and stored in liquid nitrogen.

Membranes (150-200 µg of protein / assay) are incubated at 4 °C for 120 min in 0.5 ml of a 50 mmol/l Tris-HCl buffer (pH 7.4) containing 2 mmol/l $MgCl_2$, 1 to $2 \cdot 10^{-9}$ mol/l of [3H]-2-[4-(3-azidophenyl)-2-oxo-1-pyrrolidinyl]butanamide (levetiracetam analog, a tritiated radioligand
20 currently used for the labelling of LBS), and increasing concentrations of the test substance. The non specific binding (NSB) is defined as the residual binding observed in the presence of a concentration of reference substance (e.g. 10^{-3} mol/l levetiracetam) that binds essentially all the receptors. Membrane-bound and free radioligands are separated by rapid filtration through glass fiber filters (equivalent to Whatman GF/C or GF/B; VEL, Belgium) pre-soaked in 0.1 %
25 polyethyleneimine and 10^{-3} mol/l levetiracetam to reduce non specific binding. Samples and filters are rinsed by at least 6 ml of 50 mmol/l Tris-HCl (pH 7.4) buffer. The entire filtration procedure does not exceed 10 seconds per sample. The radioactivity trapped onto the filters is counted by liquid scintillation in a β -counter (Tri-Carb 1900 or TopCount 9206, Camberra Packard, Belgium, or any other equivalent counter). Data analysis is performed by a computerized
30 non linear curve fitting method using a set of equations describing several binding models assuming populations of independent non-interacting receptors which obey to the law of mass. In tables I-A and I-B the pK_i values of various 4- or 5-substituted 2-oxo-1-piperidinyl and 5-substituted 2-oxo-1-azepanyl butanamide compounds are compared to the activity of the non-substituted levetiracetam, with a pK_i of 6.1 ± 0.1 . The lower the K_i (the higher the pK_i), the
35 stronger the binding, and thus generally the more potent the ligand (the drug).

EXAMPLE 9 : Animal model of sound-susceptible mice

The objective of this test is to evaluate the anticonvulsant potency of a compound in sound-susceptible mice, a genetic animal model with reflex seizures. In this model of primary generalised epilepsy, seizures are evoked without electrical or chemical stimulation and the seizure types are, at least in part, similar in their clinical phenomenology to seizures occurring in man (Löscher W. & Schmidt D., *Epilepsy Res.* (1998), 2, 145-181; Buchhalter J.R., *Epilepsia* (1993), 34, S31-S41).

Male or female genetically sound-sensitive mice (14-28 g; N=10), derived from a DBA strain originally selected by Dr. Lehmann of the Laboratory of Acoustic Physiology (Paris) and bred in the UCB Pharma Sector husbandry unit since 1978, were used. The experimental design consisted of several groups, one group receiving the vehicle control and the other groups different doses of the test-compound. The compounds were administered intraperitoneally 60 minutes before the induction of audiogenic seizures. The range of the doses administered had a logarithmic progression, generally between 1.0×10^{-5} mol/kg and 1.0×10^{-3} mol/kg, but lower or higher doses were tested if necessary.

For testing, the animals were placed in small cages, one mouse per cage, in a sound-attenuated chamber. After a period of orientation of 30 seconds, the acoustic stimulus (90dB, 10-20 kHz) was delivered for 30 seconds via loudspeakers positioned above each cage. During this interval, the mice were observed and the presence of the 3 phases of the seizure activity namely wild running, clonic and tonic convulsions, was recorded. The proportion of mice protected against wild running, clonic and tonic convulsions, respectively, was calculated.

For active compounds, an ED₅₀ value, i.e. the dose (expressed in moles per kg of test animal) producing 50 % protection relative to the control group, together with 95 % confidence limits, was calculated using a Probit Analysis (SAS/STAT® Software, version 6.09, PROBIT procedure) of the proportions of protected mice for each of the 3 phases of the seizure activity.

In tables I-A and I-B the ED₅₀ value (= AUD CC) for the inhibition of Clonic Convulsions in the Audiogenic seizure test are given. This test is a good animal model for the screening of anticonvulsant and/or antiepileptic compounds. Our reference compound is levetiracetam, with a ED₅₀ CC of 180 µmol/kg. The lower the ED₅₀ the more potent the compound.

In the table, the stereochemical information is contained in the two columns headed 'configuration data'. The second column indicates whether a compound has no stereogenic center (ACHIRAL), is a pure enantiomer (PURE), a racemate (RAC) or is a mixture of two or more stereoisomers, possibly in unequal proportions (MIXT). The first column contains the stereochemical assignment for each recognised center, following the IUPAC numbering used in the preceding column. A number alone indicates the existence of both configurations at that center. A number followed by 'R' or 'S' indicates the known absolute configuration at that center.

A number followed by '§' indicates the existence of only one but unknown absolute configuration at that center. The letter (A, B, C, D) in front is a way of distinguishing the various enantiomers or racemates of the same structure.

5 In the tables, the melting points are in most cases determined by the onset of the DSC curve. When a visual (fusionometer) melting point is given, the value is in parenthesis.

Table I: Synthesis intermediates

	IUPAC NAME	Configuration		Mp (°C)	LC/ MS MH ⁺	¹ H NMR
1	Methyl 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidinecarboxylate	A-1S,3§	PURE	(64-65)		
2	Methyl 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidinecarboxylate	B-1S,3§	PURE	(98-100)		
3	(2S)-2-[5-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide	A-2S,5§	PURE	(138-141)		
4	(2S)-2-[5-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide	B-2S,5§	PURE	(142-143)		
23	1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidinyl)methyl methanesulfonate	A-2S,5§	PURE			[1]
24	{1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-3-piperidinyl)methyl methanesulfonate	A-2S,5§	PURE			[2]
25	5-phenyl-2-piperidinone	A-5§	PURE			[3]
26	Ethyl 2-(2-oxo-5-phenyl-1-piperidinyl)butanoate	A-2,4§	MIXT			[4]
28	Ethyl 7-oxo-4-azepanecarboxylate	5	RAC		186	
29	Ethyl 1-[1-(tert-butoxycarbonyl)propyl]-7-oxo-4-azepanecarboxylate	2,5	MIXT		328	
30	2-[5-(ethoxycarbonyl)-2-oxo-1-azepanyl]butanoic acid	2,5	MIXT		272	
31	1-[1-(aminocarbonyl)propyl]-7-oxo-4-azepanecarboxylate	2,5	MIXT		271	
32	2-[5-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide	A-2,5	RAC		229	

Table II: ^1H NMR description

^1H NMR Number	^1H NMR Description	Solvent
[1]	0.79 (t, 3H), 1.47-1.68 (m, 2H), 1.75-1.93 (m, 2H), 2.12-2.40 (m, 3H), 3.09 (dd, 1H), 3.18 (s, 3H), 3.24 (dd, 1H), 4.20 (d, 2H), 4.84 (dd, 1H), 6.90 (broad s, 1H), 7.15 (broad s, 1H)	DMSO
[2]	0.81 (t, 3H), 1.49-1.65 (m, 2H), 1.81-1.91 (m, 2H), 2.09-2.44 (m, 3H), 3.11 (dd, 1H), 3.18 (s, 3H), 3.27-3.38 (m, 1H), 4.12 (d, 2H), 4.82 (dd, 1H), 6.92 (broad s, 1H), 7.1 (broad s, 1H)	DMSO
[3]	1.91-2.04 (m, 2H), 2.20-2.40 (m, 2H), 2.92-3.04 (m, 1H), 3.17-3.33 (m, 2H), 7.19-7.32 (m, 5H), 7.45 (broad s, 1H)	DMSO
[4]	0.87 and 0.88 (2t, 3H), 1.19 and 1.20 (2t, 3H), 1.69-2.05 (m, 4H), 2.42-2.49 (m, 2H), 3.07-3.13 (m, 1H), 3.27-3.35 (m, 2H), 4.10 (q, 1H), 4.75-4.87 (2dd, 1H), 7.23-7.35 (m, 5H)	DMSO

Table III A: 6-ring compounds

	IUPAC NAME	Configuration		Mp	PKi	AUD CC
5	(2S)-2-[5-(iodomethyl)-2-oxo-1-piperidinyl]butanamide	A-2S,5§	PURE	100.6	++	1.6E-04
6	(2S)-2-[5-(iodomethyl)-2-oxo-1-piperidinyl]butanamide	B-2S,5§	PURE	60.2	+++	2.0E-04
7	(2S)-2-[5-(azidomethyl)-2-oxo-1-piperidinyl]butanamide	A-2S,5§	PURE	70.1	+++	1.2E-04
8	(2S)-2-[5-(azidomethyl)-2-oxo-1-piperidinyl]butanamide	B-2§,5§	PURE	96.6	++	2.8E-04
9	2-(2-oxo-5-phenyl-1-piperidinyl)butanamide	A-2§,5§	PURE	167.0		2.8E-04
10	2-(2-oxo-5-phenyl-1-piperidinyl)butanamide	B-2§,5§	PURE	160.7		Inactive
11	2-(2-oxo-5-phenyl-1-piperidinyl)butanamide	C-2§,5§	PURE	158.9	+++	7.0E-05
12	2-(2-oxo-5-phenyl-1-piperidinyl)butanamide	D-2§,5§	PURE	167.8		Inactive
13	(2S)-2-[4-(iodomethyl)-2-oxo-1-piperidinyl]butanamide	A-2S,4§	PURE	144.0	+++	1.1E-04
14	(2R)-2-[4-(iodomethyl)-2-oxo-1-piperidinyl]butanamide	B-2R,4§	PURE	144.3		Inactive
15	(2S)-2-[4-(iodomethyl)-2-oxo-1-piperidinyl]butanamide	B-2S,4§	PURE	153.7	+++	2.5E-04
16	(2R)-2-[4-(iodomethyl)-2-oxo-1-piperidinyl]butanamide	A-2S,4§	PURE	153.3		Inactive

Table III B: 7-ring compounds

	IUPAC NAME	Configuration		Mp	LC/MS MH ⁺	pKi	AUD CC
17	2-[5-(iodomethyl)-2-oxo-1-azepanyl]butanamide	A-2§,5§	PURE		339	+	
18	2-[5-(iodomethyl)-2-oxo-1-azepanyl]butanamide	B-2§,5§	PURE		339	+++	1.3E-04
19	2-[5-(iodomethyl)-2-oxo-1-azepanyl]butanamide	C-2§,5§	PURE	111.7		+	Inactive
20	2-[5-(iodomethyl)-2-oxo-1-azepanyl]butanamide	D-2§,5§	PURE	111.2		++	5.6E-04

Legend

+ = pKi < 5.5

++ = 5.5 ≤ pKi < 6.5


+++ = 6.5 ≤ pKi < 7.5

++++ = pKi ≥ 7.5

inactive = inactive at 1.0E-03

CLAIMS

1. Compounds of the formula I



wherein

n represents 0 or 1 whereby R¹ is not existent when n=0 and R¹ is existent when n=1;

A¹ represents an oxygen or a sulfur atom;

X is -CONR⁷R⁸, -COOR⁹, -CO-R¹⁰ or CN;

R¹ when existent, R², R³, R⁴ and R⁵ are the same or different and each is independently hydrogen, halogen, hydroxy, thiol, amino, nitro, nitrooxy, cyano, azido, carboxy, amido, sulfonic acid, sulfonamide, alkyl, alkenyl, alkynyl, ester, ether, aryl, heterocycle, or an oxy derivative, thio derivative, amino derivative, acyl derivative, sulfonyl derivative or sulfinyl derivative,

provided that at least one of the substituents R chosen from R¹ when existent, R², R³, R⁴ or R⁵ is not hydrogen;

R⁶ is hydrogen, alkyl, aryl or -CH₂-R^{6a} wherein R^{6a} is aryl, heterocycle, halogen, hydroxy, amino, nitro or cyano;

R⁷, R⁸ and R⁹ are the same or different and each is independently hydrogen, hydroxy, alkyl, aryl, heterocycle or an oxy derivative; and

R¹⁰ is hydrogen, hydroxy, thiol, halogen, alkyl, aryl, heterocycle or a thio derivative;

their pharmaceutically acceptable salts, geometrical isomers (including cis and trans, Z and E isomers), enantiomers, diastereoisomers and mixtures thereof (including all possible mixtures of stereoisomers).

2. A compound according to claim 1 wherein n represents 0.

3. A compound according to claim 1 wherein n represents 1.

4. A compound according to claim 2 wherein R^3 and/or R^4 is different from hydrogen and R^2 and R^5 represent hydrogen.

5. A compound according to claim 3 wherein R^2 , R^3 and/or R^4 are different from hydrogen and R^1 and R^5 represent hydrogen.

6. A compound according to any of the claims 1 to 5 wherein only one R substituent chosen from R^3 or R^4 , if n represents 0, or from R^2 , R^3 or R^4 , if n represents 1, is different from hydrogen and the remaining R substituents are hydrogen.

7. A compound according to any of claims 1 to 6 wherein A^1 represents an oxygen atom.

8. A compound according to any of claims 1 to 7 wherein X represents $-\text{CONR}^7\text{R}^8$.

9. A compound according to claim 8 wherein X represents $-\text{CONH}_2$

10. A compound according to any of claims 1 to 9 wherein R^6 represents hydrogen, a C1-4 alkyl or a $\text{CH}_2\text{-R}^{6a}$ group wherein R^{6a} represents a heterocycle.

11. A compound according to claim 10 wherein R^6 represents a C1-4-alkyl, preferably ethyl.

12. A compound according to any of claims 1 to 11 wherein the carbon atom to which R^6 is attached is of the S configuration.

13. A compound according to any of the claims 1 to 12 wherein
 R^2 when $n=1$, R^3 and R^4 are the same or different and each is independently hydrogen, halogen, nitro, nitrooxy, cyano, carboxy, amido, sulfonic acid, sulfonamide, alkyl, alkenyl, alkynyl, ester, ether, aryl, heterocycle, acyl derivative, sulfonyl derivative or sulfinyl derivative;
 R^1 when existent, R^2 when $n=0$ and R^5 are hydrogen;
 R^6 is hydrogen, alkyl, aryl or $-\text{CH}_2\text{-R}^{6a}$ wherein R^{6a} is aryl, heterocycle, halogen, hydroxy, amino, nitro or cyano;

provided that, when R^6 is hydrogen, X is $-\text{CONR}^7\text{R}^8$ and that the compound is neither methyl (2R)-2-[(6R)-6-methyl-2-oxoazepanyl]-3-phenylpropanoate nor methyl (2S)-2-[(4R)-4-methyl-2-oxoazepanyl]-3-phenylpropanoate.

14. A compound according to claim 13 wherein, when R^6 is benzyl, X is $-\text{COOCH}_3$ and $n=1$, R^2 is different from methyl when R^3 and R^4 are both hydrogen and R^4 is different from methyl when R^2 and R^3 are both hydrogen.

15. A compound according to claim 13 or 14 wherein R^2 when $n=1$, R^3 and R^4 are the same or different and each is independently hydrogen; cyano; carboxy; amido;

C1-12 alkyl, each optionally substituted by one or more substituents selected from hydroxy, halogen, cyano, thiocyanato, alkoxy, azido, alkylthio, cycloalkyl, acyl, aryl and heterocycle;

C2-12 alkenyl, each optionally substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, alkyl, aryl and acyl;

C2-12 alkynyl, each optionally substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, alkyl, aryl and acyl;

acyl derivative of formula $-\text{CO}-R^{11}$, wherein R^{11} is selected from C1-12 alkyl, C2-12 alkenyl, C2-12 alkynyl, heterocycle and aryl;

ester of formula $-\text{CO}-\text{O}-R^{11}$ wherein R^{11} is selected from C1-12 alkyl, C2-12 alkenyl, C2-12 alkynyl and aryl;

heterocycle selected from triazolyl, tetrazolyl, pyrrolidinyl, pyridyl, 1-oxidopyridyl, thiomorpholinyl, benzodioxolyl, furyl, oxazolyl, pyrimidinyl, pyrrolyl, thiadiazolyl, thiazolyl, thienyl and piperazinyl, each optionally substituted by one or more substituents selected from halogen, alkyl, substituted alkyl, alkoxy, nitro, amino, acyl and phenyl;

aryl, each optionally substituted by one or more substituents selected from C1-6 alkyl, C1-6 haloalkyl, C1-6 alkoxy, C1-6 alkylthio, amino, azido, sulfonyl, aryl and nitro.

16. A compound according to claim 15, wherein R^2 when $n=1$, R^3 and R^4 are the same or different and each is independently hydrogen;

C1-7 alkyl, each optionally substituted by one or more substituents selected from hydroxy, halogen, cyano, thiocyanato, alkoxy, azido, alkylthio, cyclopropyl, acyl and phenyl;

C2-6 alkenyl, each optionally substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, cycloalkyl, phenyl and acyl;

C2-6 alkynyl, each optionally substituted by one or more substituents selected from halogen, cyano, thiocyanato, azido, alkylthio, cycloalkyl, phenyl and acyl;

heterocycle selected from tetrazolyl, pyrrolidinyl, pyridyl, furyl, pyrrolyl, thiazolyl and thienyl, each optionally substituted by one or more substituents selected from halogen, alkyl, halogen substituted alkyl, acyl, alkoxy, nitro, amino and phenyl;

phenyl, each optionally substituted by one or more substituents selected from C1-6 alkyl, C1-6 haloalkyl, halogen, C1-6 alkoxy, amino, azido, sulfonyl, phenyl and nitro.

17. A compound according to any of claims 1 to 16 wherein at least one of the R substituents
5 chosen from the group R^2 , R^3 and R^4 when $n=1$ or from the group R^3 and R^4 when $n=0$ represents independently C1-4-alkyl or C3-7-cycloalkyl, optionally substituted by one or more hydroxy, halogen, lower alkyl or/and azido.

18. A compound according to any of claims 1 to 16 wherein at least one of the R substituents
10 chosen from the group R^2 , R^3 and R^4 when $n=1$ or from the group R^3 and R^4 when $n=0$ represents independently vinyl, optionally substituted by one or more halogen or/and lower alkyl.

19. A compound according to any of claims 1 to 16 wherein at least one of the R substituents
15 chosen from the group R^2 , R^3 and R^4 when $n=1$ or from the group R^3 and R^4 when $n=0$ represents independently ethynyl, propynyl or butynyl, optionally substituted by one or more halogen or/and lower alkyl.

20. A compound according to any of claims 1 to 16 wherein at least one of the R substituents
20 chosen from the group R^2 , R^3 and R^4 when $n=1$ or from the group R^3 and R^4 when $n=0$, represents independently phenyl, optionally substituted by one or more halogen, lower alkyl, azido and/or nitro.

21. A compound according to any of claims 1 to 16 wherein at least one of the R substituent
25 chosen from the group R^2 , R^3 and R^4 when $n = 1$ or from the group R^3 and R^4 when $n = 0$, represents independently 2-or 3-thienyl, optionally substituted by one or more halogen, acyl, cyano and/or lower alkyl.

22. A compound according to any of the claims 13 to 21 wherein R^2 and R^4 when $n=1$ or R^4
30 when $n=0$ are / is hydrogen.

23. A compound according to any of the claims 13 to 21 wherein R^2 and R^3 when $n=1$ or R^3
when $n=0$ are/ is hydrogen.

24. A compound according to any of the claims 13 to 21 wherein $n=1$ and R^3 and R^4 are
35 hydrogen.

25. A compound according to claim 1 selected from the group consisting of 2-[5-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-5-propyl-1-piperidinyl)butanamide, 2-[2-oxo-5-(3,3,3-trifluoropropyl)-1-piperidinyl]butanamide, 2-[5-(cyclopropylmethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(iodomethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(azidomethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-5-phenyl-1-piperidinyl)butanamide, 2-[2-oxo-5-(2-thienyl)-1-piperidinyl]butanamide, 2-[2-oxo-5-(3-thienyl)-1-piperidinyl]butanamide, 2-[5-(3-chlorophenyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(3-azidophenyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2,2-difluorovinyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2,2-dibromovinyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2,2-dichlorovinyl)-2-oxo-1-piperidinyl]butanamide, 2-(5-ethynyl-2-oxo-1-piperidinyl)butanamide, 2-[5-(5-methyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(5-formyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(5-cyano-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(3-bromo-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(4-methyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[2-oxo-5-(3,3,3-trifluoro-1-propynyl)-1-piperidinyl]butanamide, 2-[2-oxo-5-(1-propynyl)-1-piperidinyl]butanamide, 2-[5-(cyclopropylethynyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(3-methyl-1-butynyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(1-butynyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2,2-difluoropropyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2-chloro-2,2-difluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(2-bromo-2,2-difluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-4-propyl-1-piperidinyl)butanamide, 2-[2-oxo-4-(3,3,3-trifluoropropyl)-1-piperidinyl]butanamide, 2-[4-(cyclopropylmethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(iodomethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(azidomethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-4-phenyl-1-piperidinyl)butanamide, 2-[2-oxo-4-(2-thienyl)-1-piperidinyl]butanamide, 2-[2-oxo-4-(3-thienyl)-1-piperidinyl]butanamide, 2-[4-(3-chlorophenyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(3-azidophenyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2-difluorovinyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2-dibromovinyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2-dichlorovinyl)-2-oxo-1-piperidinyl]butanamide, 2-(4-ethynyl-2-oxo-1-piperidinyl)butanamide, 2-[4-(5-methyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(5-formyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(5-cyano-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(3-bromo-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(4-methyl-2-thienyl)-2-oxo-1-piperidinyl]butanamide, 2-[2-oxo-4-(3,3,3-trifluoro-1-propynyl)-1-piperidinyl]butanamide, 2-[2-oxo-4-(1-propynyl)-1-piperidinyl]butanamide, 2-[4-(cyclopropylethynyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(3-methyl-1-butynyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(1-butynyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2-difluoropropyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2-chloro-2,2-difluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-

(2-bromo-2,2-difluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[4-(2,2,2-trifluoroethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-5-propyl-1-azepanyl]butanamide, 2-[2-oxo-5-(3,3,3-trifluoropropyl)-1-azepanyl]butanamide, 2-[5-(cyclopropylmethyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(iodomethyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(azidomethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-5-phenyl-1-azepanyl]butanamide, 2-[2-oxo-5-(2-thienyl)-1-azepanyl]butanamide, 2-[2-oxo-5-(3-thienyl)-1-azepanyl]butanamide, 2-[5-(3-chlorophenyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(3-azidophenyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2-difluorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2-dibromovinyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2-dichlorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[5-ethynyl-2-oxo-1-azepanyl]butanamide, 2-[5-(5-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(5-formyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(5-cyano-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(3-bromo-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(4-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[2-oxo-5-(3,3,3-trifluoro-1-propynyl)-1-azepanyl]butanamide, 2-[2-oxo-5-(1-propynyl)-1-azepanyl]butanamide, 2-[5-(cyclopropylethynyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(3-methyl-1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2-difluoropropyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2-chloro-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2-bromo-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[5-(2,2,2-trifluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-6-propyl-1-azepanyl]butanamide, 2-[2-oxo-6-(3,3,3-trifluoropropyl)-1-azepanyl]butanamide, 2-[6-(cyclopropylmethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(iodomethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(azidomethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-6-phenyl-1-azepanyl]butanamide, 2-[2-oxo-6-(2-thienyl)-1-azepanyl]butanamide, 2-[2-oxo-6-(3-thienyl)-1-azepanyl]butanamide, 2-[6-(3-chlorophenyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(3-azidophenyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2-difluorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2-dibromovinyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2-dichlorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[6-ethynyl-2-oxo-1-azepanyl]butanamide, 2-[6-(5-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(5-formyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(5-cyano-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(3-bromo-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(4-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[2-oxo-6-(3,3,3-trifluoro-1-propynyl)-1-azepanyl]butanamide, 2-[2-oxo-6-(1-propynyl)-1-azepanyl]butanamide, 2-[6-(cyclopropylethynyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(3-methyl-1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(1-butynyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2-difluoropropyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2-chloro-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2-bromo-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[6-(2,2,2-trifluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-4-propyl-1-

azepanyl]butanamide, 2-[2-oxo-4-(3,3,3-trifluoropropyl)-1-azepanyl]butanamide, 2-[4-(cyclopropylmethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(iodomethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(azidomethyl)-2-oxo-1-azepanyl]butanamide, 2-(2-oxo-4-phenyl-1-azepanyl]butanamide, 2-[2-oxo-4-(2-thienyl)-1-azepanyl]butanamide, 2-[2-oxo-4-(3-thienyl)-1-azepanyl]butanamide, 2-[4-(3-chlorophenyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(3-azidophenyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2-difluorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2-dibromovinyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2-dichlorovinyl)-2-oxo-1-azepanyl]butanamide, 2-[4-ethynyl-2-oxo-1-azepanyl]butanamide, 2-[4-(5-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(5-formyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(5-cyano-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(3-bromo-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(4-methyl-2-thienyl)-2-oxo-1-azepanyl]butanamide, 2-[2-oxo-4-(3,3,3-trifluoro-1-propynyl)-1-azepanyl]butanamide, 2-[2-oxo-4-(1-propynyl)-1-azepanyl]butanamide, 2-[4-(cyclopropylethynyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(3-methyl-1-butyryl)-2-oxo-1-azepanyl]butanamide, 2-[4-(1-butyryl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2-difluoropropyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2-chloro-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2-bromo-2,2-difluoroethyl)-2-oxo-1-azepanyl]butanamide, 2-[4-(2,2,2-trifluoroethyl)-2-oxo-1-azepanyl]butanamide.

26. A compound according to claim 1 selected from the group consisting of (2S)-2-[5-(iodomethyl)-2-oxo-1-piperidinyl]butanamide, (2S)-2-[5-(azidomethyl)-2-oxo-1-piperidinyl]butanamide, 2-(2-oxo-5-phenyl-1-piperidinyl]butanamide, (2S)-2-[4-(iodomethyl)-2-oxo-1-piperidinyl]butanamide, 2-[5-(iodomethyl)-2-oxo-1-azepanyl]butanamide.

27. A pharmaceutical composition comprising as active ingredient a therapeutically effective amount of a compound according to any of claims 1 to 26 and a pharmaceutically acceptable adjuvant, diluent or carrier.

28. A compound according to any of claims 1 to 26 for use as a medicine.

29. Use of a compound according to any of claims 1 to 26, in the manufacture of a medicament for the treatment of epilepsy, epileptogenesis, seizure disorders, convulsions and other neurological disorders including bipolar disorders, mania, depression, anxiety, migraine, trigeminal and other neuralgia, chronic pain, neuropathic pain, cerebral ischemia, cardiac arrhythmia, myotonia, cocaine abuse, stroke, myoclonus, essential tremor, dyskinesia and other movement disorders, neonatal cerebral haemorrhage, amyotrophic lateral sclerosis, spasticity, Parkinson's disease and other degenerative diseases, bronchial asthma, asthmatic status and


allergic bronchitis, asthmatic syndrome, bronchial hyperreactivity and bronchospastic syndromes as well as allergic and vasomotor rhinitis and rhinoconjunctivitis.

30. Use according to claim 29 in the manufacture of a medicament for the treatment of
5 epilepsy, epileptogenesis, seizure disorders and convulsions.


31. A method for treating epilepsy, epileptogenesis, seizure disorders, convulsions and other neurological disorders including bipolar disorders, mania, depression, anxiety, migraine, trigeminal and other neuralgia, chronic pain, neuropathic pain, cerebral ischemia, cardiac
10 arrhythmia, myotonia, cocaine abuse, stroke, myoclonus, essential tremor, dyskinesia and other movement disorders, neonatal cerebral haemorrhage, amyotrophic lateral sclerosis, spasticity, Parkinson's disease and other degenerative diseases, bronchial asthma, asthmatic status and allergic bronchitis, asthmatic syndrome, bronchial hyperreactivity and bronchospastic syndromes as well as allergic and vasomotor rhinitis and rhinoconjunctivitis, in a mammal in
15 need of such treatment, comprising administering a therapeutic dose of at least one compound according to any of claims 1 to 26.

32. A method according to claim 31 for treating epilepsy, epileptogenesis, seizure disorders and convulsions.


20 33. A process for preparing a compound of general formula (I) of claim 1 wherein A^1 represents O and X represents $-\text{CONR}^7\text{R}^8$ or $-\text{COOR}^9$, wherein this process comprises the cyclisation of an aminoester of formula (A) wherein Q^1 represents a linear or branched C1-4-alkyl, and wherein said aminoester of formula (A) wherein n represents 0 and R^3 represents $-\text{COOR}^{13}$, or wherein n represents 1 and R^2 represents $-\text{COOR}^{13}$, wherein R^{13} represents a linear or branched alkyl group optionally optically active, are obtained by reaction of a compound of formula (B) with an unsaturated diester derivative of formula (C) or (D), respectively, according to the following scheme:




34. A process for preparing a compound of general formula (I) of claim 1 wherein A^1 represents O, X represents $-CONR^7R^8$ and n represents 0, wherein this process comprises the reaction of a compound of formula (E), wherein X^1 represents a halogen atom, preferably an iodine or a chlorine atom, X^2 represents $-OQ^1$ or a halogen atom, wherein Q^1 represents a linear or branched C1-4-alkyl, with a compound of formula (F); whereby said compound of formula (E) is obtained by opening of a lactone of formula (G) in the presence of a halogenating agent, and said compound of formula (G) is prepared by reaction of a compound of formula (J) with an organolithium compound of formula R^4Li in the presence of CuI or, when in formula (G) $R^2=H$, by reduction of a carboxylate of formula (H) wherein Q^4 =methyl or ethyl and in the presence of a borohydride reagent, all reactions according to the following scheme :



35. A process for preparing a compound of general formula (I) of claim 1 wherein A¹ represents O and X represents -COOR⁹, R⁹ representing an alkyl group, wherein this process
 5 comprises the reaction of a compound of formula (K) wherein M represents an alkali metal, with a compound of formula (L) wherein X³ represents a halogen atom, according to the following scheme:



36. A process for preparing a compound of general formula (I) of claim 1 wherein A¹ represents O and X represents -CONR⁷R⁸ and none of the substituents R¹, R², R³, R⁴, R⁵ and R⁶ being substituted by carboxylic or sulfonic acids or esters, or ketone, wherein this process
 15 comprises the ammonolysis of the corresponding ester of formula (II) wherein R⁹ represents a hydrogen atom or a linear or branched C1-4-alkyl group.



37. Intermediates prepared by any of the processes according to claims 33 to 36.

- 5 38. Intermediates according to claim 37 selected from the group consisting of methyl 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-(3S)-3-piperidinecarboxylate and methyl 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-(3R)-3-piperidinecarboxylate and their mixtures, (2S)-2-[(5R)-5-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide and (2S)-2-[(5S)-5-(hydroxymethyl)-2-oxo-1-piperidinyl]butanamide and their mixtures, 1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-(3R)-3-piperidinyl)methyl methanesulfonate and {1-[(1S)-1-(aminocarbonyl)propyl]-6-oxo-(3S)-3-piperidinyl)methyl methanesulfonate and their mixtures, 5-phenyl-2-piperidinone, ethyl 2-(2-oxo-5-phenyl-1-piperidinyl)butanoate, ethyl 7-oxo-4-azepanecarboxylate, ethyl 1-[1-(tert-butoxycarbonyl)propyl]-7-oxo-4-azepanecarboxylate, 2-[5-(ethoxycarbonyl)-2-oxo-1-azepanyl]butanoic acid, 1-[1-(aminocarbonyl)propyl]-7-oxo-4-azepanecarboxylate, 2-[5-(hydroxymethyl)-2-oxo-1-azepanyl]butanamide.
- 10
- 15

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/05503

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D211/76 C07D211/78 C07D409/04 C07D223/10 A61K31/45
A61K31/55 A61P25/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PIRO, J. ET AL: "Solid phase synthesis of enantiomerically pure polyhydroxyvalerolactams" TETRAHEDRON LETT. (2001), 42(5), 871-873, XP002180740 examples 1,2A,2B --- -/--	1

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

19 August 2002

Date of mailing of the international search report

28/08/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seitner, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/05503

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ESTIARTE M A ET AL: "Synthesis and Synthetic Applications of 3-Amino-DELTA-piperidein-2 -ones: Synthesis of Methionine-derived Pseudopeptides" TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 55, no. 33, 13 August 1999 (1999-08-13), pages 10173-10186, XP004174815 ISSN: 0040-4020 examples 20,21	1
A	* scheme 4 *	34
X	US 5 776 950 A (DUTTON FRED E ET AL) 7 July 1998 (1998-07-07) examples 5,6,8	1
X	DUTTON F E ET AL: "Epsilon-lactam Analogs of the Anthelmintic Cyclopeptide PF1022A" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 39, no. 30, 23 July 1998 (1998-07-23), pages 5313-5316, XP004123220 ISSN: 0040-4039 example 11	1
X	US 5 719 296 A (JONES A BRIAN ET AL) 17 February 1998 (1998-02-17) examples 50,59,60 claims 8,9	1,27
X	EP 0 783 002 A (SQUIBB & SONS INC) 9 July 1997 (1997-07-09) page 55, line 48 examples 44B,55	1
X	ROBL J A ET AL: "DUAL METALLOPROTEASE INHIBITORS. 6. INCORPORATION OF BICYCLIC AND SUBSTITUTED MONOCYCLIC AZEPINONES AS DIPEPTIDE SURROGATES IN ANGIOTENSIN-CONVERTING ENZYME/NEUTRAL ENDOPEPTIDASE INHIBITORS" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 39, no. 2, 19 January 1996 (1996-01-19), pages 494-502, XP000749701 ISSN: 0022-2623 examples 5A-5Q,12A,12B abstract	1,27
	--- -/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/05503

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ROBL, JEFFREY A. ET AL: "Peptidomimetic Synthesis: A Novel, Highly Stereoselective Route to Substituted Freidinger Lactams" J. AM. CHEM. SOC. (1994), 116(6), 2348-55, XP002180741 examples 16B, 16E, 17B, 17E	1
A	* equation (1) *	35
X	DEPRIEST, SCOTT A. ET AL: "3D-QSAR of angiotensin-converting enzyme and thermolysin inhibitors: a comparison of CoMFA models based on deduced and experimentally determined active site geometries" J. AM. CHEM. SOC. (1993), 115(13), 5372-84, XP002180742 abstract; example 44	1, 27
X	DE LASZLO, S. E. ET AL: "Synthesis and use of 3-amino-4-phenyl-2-piperidones and 4-amino-2-benzazepin-3-ones as conformationally restricted phenylalanine isosteres in renin inhibitors" J. MED. CHEM. (1992), 35(5), 833-46, XP002180743 abstract examples 12B, 14B, 12A, 14A	1, 27
A	* scheme II *	34
X	EP 0 365 992 A (ABBOTT LAB) 2 May 1990 (1990-05-02) page 52; example 25	1
X	AEBI, JOHANNES D. ET AL: "Synthesis, conformation and immunosuppressive activity of a conformationally restricted cyclosporin lactam analog" J. MED. CHEM. (1988), 31(9), 1805-15, XP002180744 example 20	1
X	EP 0 046 291 A (MERCK & CO INC) 24 February 1982 (1982-02-24) claims 8, 10 * formula V, VI, XVIII *	1

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/05503

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MILEWSKA M J ET AL: "Synthesis of Lactams by Regioselective Reduction of Cyclic Dicarboximides" SYNTHESIS, vol. 12, - 1996 pages 1485-1488, XP002180745 example 3H ---	38
Y	WO 95 35308 A (VERTEX PHARMA) 28 December 1995 (1995-12-28) page 117, line 6 - line 12; claims 74,76 example 92 ---	1,27,29
Y	GB 1 309 692 A (UCB SA) 14 March 1973 (1973-03-14) page 1, line 19 - line 20 page 2, line 27 - line 32 examples 1-5 claims 1,40 ---	1,27,29
Y	US 5 334 720 A (SCHMIESING RICHARD J ET AL) 2 August 1994 (1994-08-02) column 1, paragraph 2 example 1 claims 1,4 ---	1,27,29
Y	PROUS: "LEVETIRACETAM" DRUGS OF THE FUTURE, BARCELONA, ES, vol. 19, no. 2, 1994, pages 111-113, XP000908882 ISSN: 0377-8282 the whole document ---	1,27,29
A	ANGELUCCI, LUCIANO ET AL: "Synthesis and amnesia-reversal activity of a series of 7- and 5-membered 3-acylamino lactams" J. MED. CHEM. (1993), 36(11), 1511-19, XP002180746 abstract table II * scheme II * examples 23,24,26,27 ---	1,27,29, 35
A	UMA SREENIVASAN ET AL: "Synthesis and dopamine receptor modulating activity of lactam conformationally constrained analogues of Pro-Leu-Gly-NH ₂ " JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 36, no. 2, 1993, pages 256-263, XP002123346 ISSN: 0022-2623 abstract * scheme I * examples 6-8,13,14,16,17,19,20 ---	1,27,29, 35
	--- -/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/05503

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 19795 A (DIMENSIONAL PHARM INC ;MARUGAN SANCHEZ JUAN JOSE (US); HALL JONATH) 22 March 2001 (2001-03-22) page 14, line 16 - line 18 example 1 ---	1,27,29
A	MICOUIN L ET AL: "Asymmetric Synthesis. XXXIX. Synthesis of 3-substituted Piperidin-2-ones from Chiral Non-Racemic Lactams" TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 52, no. 22, 27 May 1996 (1996-05-27), pages 7719-7726, XP004106225 ISSN: 0040-4020 example 1 -----	35

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 02/05503

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 31 and 32 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-13 (all partially); 14 (completely); 15-24 (all partially); 27-37 (all partially)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT).

For these reasons, it appears impossible to execute a meaningful search and /or to issue a complete search report over the whole breadth of above mentioned claims.

The search and the report for those claims can only be considered complete for compounds of formula (I) of claim 1 for which:

-R6 represents unsubstituted, straight or branched alkyl,

-oxy-derivative means -O-R11,

-thio-derivative means -S-R11,

-amino-derivative means -NH-R11 or -NR11R11.

In above mentioned definition, R11 represents unsubstituted alkyl, unsubstituted alkenyl, unsubstituted alkynyl, or unsubstituted aryl.

The search for claim 37 was restricted to intermediates according to claims 23-26, which contributes to the structural differentiation of the end-products of formula (I) over the state of the art and which comprises the moiety -CH(R6)-X, namely to compounds of formula (A) of claim 33 and compounds of formula (II) of claim 36.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/05503

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5776950	A	07-07-1998	AU 6860296 A	27-03-1997
			BR 9610514 A	30-03-1999
			CA 2228632 A1	13-03-1997
			CN 1196056 A	14-10-1998
			EP 0863907 A1	16-09-1998
			JP 11513361 T	16-11-1999
			WO 9709331 A1	13-03-1997
US 5719296	A	17-02-1998	AU 7604196 A	22-05-1997
			WO 9716425 A1	09-05-1997
EP 0783002	A	09-07-1997	EP 0783002 A2	09-07-1997
			AT 162800 T	15-02-1998
			AU 674629 B2	02-01-1997
			AU 2852595 A	19-10-1995
			AU 3862793 A	25-11-1993
			BR 1101081 A3	28-03-2000
			CA 2096460 A1	19-11-1993
			CN 1084849 A , B	06-04-1994
			CZ 9300936 A3	15-12-1993
			DE 69316717 D1	05-03-1998
			DE 69316717 T2	27-08-1998
			DK 599444 T3	21-09-1998
			EP 0599444 A1	01-06-1994
			ES 2111712 T3	16-03-1998
			FI 932249 A	19-11-1993
			GR 3026665 T3	31-07-1998
			HK 1003597 A1	30-10-1998
			HU 68391 A2	28-06-1995
			HU 218482 B	28-09-2000
			IL 105729 A	04-01-1998
			JP 6056790 A	01-03-1994
			KR 277453 B1	15-12-2000
			MX 9302876 A1	01-12-1993
			NO 931795 A	19-11-1993
			NZ 247642 A	26-09-1995
			PL 298988 A1	24-01-1994
			RU 2124503 C1	10-01-1999
			SG 54138 A1	16-11-1998
			SK 49993 A3	10-08-1994
			US 5552397 A	03-09-1996
			US 5723602 A	03-03-1998
			US 5723457 A	03-03-1998
			US 5859239 A	12-01-1999
			ZA 9303461 A	14-12-1993
EP 0365992	A	02-05-1990	AU 4849390 A	28-05-1990
			CA 2000929 A1	19-04-1990
			DK 70491 A	17-06-1991
			EP 0365992 A1	02-05-1990
			EP 0439556 A1	07-08-1991
			JP 4501566 T	19-03-1992
			NZ 231053 A	26-02-1991
			WO 9004917 A1	17-05-1990
			US 5153358 A	06-10-1992
			US 5164388 A	17-11-1992
			US 5268476 A	07-12-1993

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/05503

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0046291	A	24-02-1982	DE 3173304 D1	06-02-1986
			DK 364181 A	19-02-1982
			EP 0046291 A2	24-02-1982
			ES 504791 D0	01-04-1983
			ES 8305337 A1	01-07-1983
			ES 519057 D0	16-03-1984
			ES 8403460 A1	16-06-1984
			GR 74635 A1	29-06-1984
			JP 1635513 C	20-01-1992
			JP 2054342 B	21-11-1990
			JP 57112373 A	13-07-1982
			PT 73537 A ,B	01-09-1981
			US 4629787 A	16-12-1986
			US 4680392 A	14-07-1987
			FI 813378 A	14-01-1983
			NO 813627 A	14-01-1983
WO 9535308	A	28-12-1995	US 5756466 A	26-05-1998
			US 5656627 A	12-08-1997
			US 5847135 A	08-12-1998
			AP 797 A	07-01-2000
			AU 709114 B2	19-08-1999
			AU 2944695 A	15-01-1996
			BG 101130 A	29-08-1997
			BR 9508051 A	21-10-1997
			CA 2192089 A1	28-12-1995
			CN 1159196 A	10-09-1997
			CZ 9603698 A3	11-06-1997
			EP 0784628 A1	23-07-1997
			FI 965036 A	14-02-1997
			HU 76622 A2	28-10-1997
			JP 10504285 T	28-04-1998
			NO 965365 A	17-02-1997
			NZ 289560 A	29-09-1999
			PL 318220 A1	26-05-1997
			SK 160996 A3	10-09-1997
			WO 9535308 A1	28-12-1995
			US 6025147 A	15-02-2000
			US 5716929 A	10-02-1998
			US 6103711 A	15-08-2000
			US 5973111 A	26-10-1999
			US 2002099042 A1	25-07-2002
			ZA 9504988 A	17-12-1996
GB 1309692	A	14-03-1973	BE 762728 A4	10-08-1971
			CH 530395 A	15-11-1972
			CS 167911 B2	28-05-1976
			CS 167912 B2	28-05-1976
			DE 2106418 A1	19-08-1971
			DK 137451 B	06-03-1978
			ES 388135 A2	01-05-1973
			FI 54916 B	29-12-1978
			FR 2081508 A6	03-12-1971
			JP 53005301 B	25-02-1978
			JP 986288 C	07-02-1980
			JP 53130656 A	14-11-1978
			JP 54020491 B	23-07-1979
			MY 7274 A	31-12-1974

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/05503

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1309692	A		NL 7101697 A ,B,	17-08-1971
			SU 508185 A3	25-03-1976
			SU 508186 A3	25-03-1976
			YU 30671 A ,B	31-10-1979
			YU 144578 A1	31-10-1982
			ZA 7100877 A	27-10-1971
US 5334720	A	02-08-1994	WO 9523135 A1	31-08-1995
			AU 6113594 A	11-09-1995
WO 0119795	A	22-03-2001	AU 7480800 A	17-04-2001
			BR 0013926 A	23-07-2002
			EP 1212300 A1	12-06-2002
			NO 20021224 A	13-05-2002
			WO 0119795 A1	22-03-2001